arXiv:1409.1453v1 [math.RA] 4 Sep 2014

A simultaneous decomposition of seven matrices over real quaternion algebra and its applications[1]

Zhuo-Heng He, Qing-Wen Wang*

Department of Mathematics, Shanghai University, Shanghai 200444. P.R. China

E-mail: wqw@staff.shu.edu.cn (Q.W. Wang), hzh19871126@126.com (Z.H. He)

Abstract: Let $\mathbb{H}$ be the real quaternion algebra and $\mathbb{H}^{n\times m}$ denote the set of all $n\times m$ matrices over $\mathbb{H}$. In this paper, we construct a simultaneous decomposition of seven general real quaternion matrices with compatible sizes: $A\in\mathbb{H}^{m\times n}, B\in\mathbb{H}^{m\times p_1}, C\in\mathbb{H}^{m\times p_2}, D\in\mathbb{H}^{m\times p_3}, E\in\mathbb{H}^{q_1\times n}, F\in\mathbb{H}^{q_2\times n}, G\in\mathbb{H}^{q_3\times n}$. As applications of the simultaneous matrix decomposition, we give solvability conditions, general solutions, as well as the range of ranks of the general solutions to the following two real quaternion matrix equations $BXE+CYF+DZG=A$ and $BX+WE+CYF+DZG=A$, where A,B,C,D,E,F, and G are given real quaternion matrices.

Keywords: Quaternion; Division ring; Matrix decomposition; Matrix equation; Rank; General solution

2010 AMS Subject Classifications: 15A21, 15A22, 15A24, 15A33, 15A03

1. Introduction

Throughout this paper, let $\mathbb{R},\mathbb{C}$, and $\mathbb{H}^{m\times n}$ stand, respectively, for the real number field, the complex number field, and the set of all $m\times n$ matrices over the real quaternion algebra

$$\mathbb{H}=\big\{a_0+a_1i+a_2j+a_3k\big|\ i^2=j^2=k^2=ijk=-1, a_0,a_1,a_2,a_3\in\mathbb{R}\big\}.$$

The rank of a quaternion matrix A is defined to be the maximum number of columns of A which are right linearly independent [26]. It is easy to see that for any nonsingular matrices P and Q of appropriate sizes, A and PAQ have the same rank [26]. Moreover, for $A\in\mathbb{H}^{m\times n}$, by [10], there exist invertible matrices P and Q such that

$$PAQ=\begin{pmatrix} I_r & 0\\ 0 & 0\end{pmatrix}$$

[1]This research was supported by the grants from the National Natural Science Foundation of China (11171205), the Natural Science Foundation of Shanghai (11ZR1412500), the Key Project of Scientific Research Innovation Foundation of Shanghai Municipal Education Commission (13ZZ080).

* Corresponding author

where $r = r(A)$, I_r is the $r \times r$ identity matrix. The rank of a block real quaternion matrix

$$\begin{pmatrix} A_{11} & A_{12} & \cdots & A_{1,n} \\ A_{21} & A_{22} & \cdots & A_{2,n} \\ \vdots & \vdots & \ddots & \vdots \\ A_{m,1} & A_{m,2} & \cdots & A_{m,n} \end{pmatrix}$$

is denoted by $r_{a_{11}a_{12}\cdots a_{1,n}|a_{21}a_{22}\cdots a_{2,n}|\cdots|a_{m,1}a_{m,2}\cdots a_{m,n}}$. If the block real quaternion matrix has zero, we use "0" instead of zero in the subscript. For instance, the ranks of the following block real quaternion matrices

$$(B,\ C,\ D),\ \begin{pmatrix} E \\ F \\ G \end{pmatrix},\ \begin{pmatrix} D & 0 & A & B & 0 \\ D & A & 0 & 0 & C \\ 0 & E & 0 & 0 & 0 \\ 0 & 0 & F & 0 & 0 \\ 0 & F & 0 & 0 & 0 \end{pmatrix}$$

are represented by r_{bcd}, $r_{e|f|g}$, $r_{d0ab0|da00c|0e000|00f00|0f000}$, respectively. The set of all $n \times n$ invertible real quaternion matrices is denoted by $GL_n(\mathbb{H})$.

Quaternions were introduced by Irish mathematician Sir William Rowan Hamilton in 1843. It is well known that $\mathbb{H}$ is an associative and noncommutative division algebra. General properties of quaternion and real quaternion matrices can be found in [26]. Nowadays real quaternion matrices have always been at the heart of computer science, quantum physics, signal and color image processing, and so on (e.g. [1], [2], [11], [15]-[18]).

In mathematics, engineering, signal, circuit and others, many problems can be transformed into the decomposition of multiple matrices (e.g. [3], [14], [25]). In 1981, Paige and Saunders [12] introduced the generalized singular value decomposition of two matrices with the same row number. In 1991, Zha [25] gave a restricted singular values of a general matrix triplet $\left(\begin{smallmatrix} A & B \\ C & \end{smallmatrix}\right)$ over $\mathbb{C}$. In 1991, Bart De Moor and G.H. Golub [4] derived a generalization of the OSVD. Several applications were discussed in [4]. Moreover, Bart De Moor and Paul Van Dooren [5] presented the generalized singular value decompositions for k general matrices $(A_1, A_2, \ldots, A_k)$, where $A_1 \in \mathbb{C}^{n_0 \times n_1}, A_2 \in \mathbb{C}^{n_1 \times n_2}, \ldots, A_k \in \mathbb{C}^{n_{k-1} \times n_k}$. In 1994, C.C. Paige and M.S. Wei [13] introduced the history and generality of the CS Decomposition. In 2000, Delin Chu, Lieven De Lathauwer and Bart De Moor [6] proved that the restricted singular value decomposition of a general matrix triplet $\left(\begin{smallmatrix} A & B \\ C & \end{smallmatrix}\right)$ can be computed using a CSD-based QR-type method. In 2011, Wang, van der Woude and Yu [19] derived the decomposition of three general matrices with the same row number over an arbitrary division ring. In 2012, Wang, Zhang and van der Woude [20] gave a simultaneous decomposition concerning the general matrix quaternity $\left(\begin{smallmatrix} A & B & C \\ D & & \end{smallmatrix}\right)$ over an arbitrary division ring. Quite recently, He and Wang [9] constructed a simultaneous decomposition of five general real quaternion matrices $\left(\begin{smallmatrix} A & B & C \\ D & & \\ E & & \end{smallmatrix}\right)$. He and Wang [9] gave the rang of ranks of the real quaternion matrix expression $A - BXD - CYE$ by using the simultaneous decomposition of five general real quaternion matrices $\left(\begin{smallmatrix} A & B & C \\ D & & \\ E & & \end{smallmatrix}\right)$.

The remainder of the paper is organized as follows. In Section 2, we propose the simultaneous decomposition of the matrix array

$$\begin{pmatrix} A & B & C & D \\ E & & & \\ F & & & \\ G & & & \end{pmatrix}, \tag{1.1}$$

where $A \in \mathbb{H}^{m\times n}, B \in \mathbb{H}^{m\times p_1}, C \in \mathbb{H}^{m\times p_2}, D \in \mathbb{H}^{m\times p_3}, E \in \mathbb{H}^{q_1\times n}, F \in \mathbb{H}^{q_2\times n}$ and $G \in \mathbb{H}^{q_3\times n}$ are general real quaternion matrices. In Section 3, we discuss several applications of the simultaneous decomposition. In Section 3.2, we derive solvability conditions and the general solution to the real quaternion matrix equation

$$BXE + CYF + DZG = A. \tag{1.2}$$

In Section 3.3, we give the range of ranks of the general solution to the real quaternion matrix equation (1.2). In Section 3.4, we present solvability conditions and the general solution to the real quaternion matrix equation

$$BX + WE + CYF + DZG = A. \tag{1.3}$$

In Section 3.5, we derive the range of ranks of the general solution to the real quaternion matrix equation (1.3).

2. A simultaneous decomposition of the general matrix array (1.1) over $\mathbb{H}$

We begin with the following lemma that is a basic tool for obtaining the main result.

Lemma 2.1. [19] *Let $B \in \mathbb{H}^{m\times p_1}, C \in \mathbb{H}^{m\times p_2}$ and $D \in \mathbb{H}^{m\times p_3}$ be given. Then there exist $\widetilde{P} \in GL_m(\mathbb{H}), \widetilde{T_1} \in GL_{p_1}(\mathbb{H}), \widetilde{T_2} \in GL_{p_2}(\mathbb{H})$ and $\widetilde{T_3} \in GL_{p_3}(\mathbb{H})$ such that*

$$B = \widetilde{P}\widetilde{S_B}\widetilde{T_1}, \qquad C = \widetilde{P}\widetilde{S_C}\widetilde{T_2}, \qquad D = \widetilde{P}\widetilde{S_D}\widetilde{T_3}, \tag{2.1}$$

where

$$(\widetilde{S_B}, \widetilde{S_C}, \widetilde{S_D}) = \begin{pmatrix} I & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & I & 0 & 0 & 0 & 0 & 0 & 0 & I & 0 & m_1 \\ 0 & I & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & I & 0 & 0 & 0 & 0 & 0 & 0 & 0 & m_2 \\ 0 & 0 & I & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & I & 0 & 0 & m_3 \\ 0 & 0 & 0 & I & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & I & 0 & 0 & 0 & 0 & m_4 \\ 0 & 0 & 0 & 0 & I & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & m_5 \\ 0 & 0 & 0 & 0 & 0 & 0 & I & 0 & 0 & 0 & 0 & 0 & 0 & I & 0 & 0 & 0 & 0 & m_4 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & I & 0 & 0 & 0 & 0 & 0 & 0 & I & 0 & 0 & 0 & m_6 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & I & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & m_7 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & I & 0 & 0 & 0 & 0 & 0 & m_8 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & m - r_{bcd} \end{pmatrix},$$

where

$$m_4 + m_6 + m_7 = r_{bc} - r_b,\ m_1 + m_2 = r_b + r_c - r_{bc},\ m_8 = r_{bcd} - r_{bc},$$

$$m_4 + m_6 = r_{bc} + r_{bd} - r_{bcd} - r_b,\ m_1 + m_3 = r_b + r_d - r_{bd},\ m_3 + m_4 = r_{bc} + r_{cd} - r_{bcd} - r_c.$$

Now we give the main theorem of this section.

Theorem 2.2. *Let $A \in \mathbb{H}^{m\times n}, B \in \mathbb{H}^{m\times p_1}, C \in \mathbb{H}^{m\times p_2}, D \in \mathbb{H}^{m\times p_3}, E \in \mathbb{H}^{q_1\times n}, F \in \mathbb{H}^{q_2\times n}$ and $G \in \mathbb{H}^{q_3\times n}$ be given. Then there exist $P \in GL_m(\mathbb{H}),\ Q \in GL_n(\mathbb{H}),\ T_1 \in GL_{p_1}(\mathbb{H}),\ T_2 \in GL_{p_2}(\mathbb{H}),\ T_3 \in GL_{p_3}(\mathbb{H}),\ V_1 \in GL_{q_1}(\mathbb{H}),\ V_2 \in GL_{q_2}(\mathbb{H})$, and $V_3 \in GL_{q_3}(\mathbb{H})$ such that*

$$A = PS_AQ, B = PS_BT_1, C = PS_CT_2, D = PS_DT_3, E = V_1S_EQ, F = V_2S_FQ, G = V_3S_GQ, \tag{2.2}$$

where

$$S_A = \begin{pmatrix} A_{11} & \cdots & A_{19} & A_{1,10} & 0 \\ \vdots & \ddots & \vdots & \vdots & \vdots \\ A_{91} & \cdots & A_{99} & A_{9,10} & 0 \\ A_{10,1} & \cdots & A_{10,9} & 0 & 0 \\ 0 & \cdots & 0 & 0 & I_t \end{pmatrix}, \tag{2.3}$$

$$(S_B, S_C, S_D) = \begin{pmatrix} I & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & I & 0 & 0 & 0 & 0 & 0 & 0 & I & 0 & m_1 \\ 0 & I & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & I & 0 & 0 & 0 & 0 & 0 & 0 & 0 & m_2 \\ 0 & 0 & I & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & I & 0 & 0 & m_3 \\ 0 & 0 & 0 & I & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & I & 0 & 0 & 0 & 0 & m_4 \\ 0 & 0 & 0 & 0 & I & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & m_5 \\ 0 & 0 & 0 & 0 & 0 & 0 & I & 0 & 0 & 0 & 0 & 0 & 0 & I & 0 & 0 & 0 & 0 & m_4 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & I & 0 & 0 & 0 & 0 & 0 & 0 & I & 0 & 0 & 0 & m_6 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & I & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & m_7 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & I & 0 & 0 & 0 & 0 & 0 & m_8 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & m - r_{bcd} - t \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & t \end{pmatrix}, \tag{2.4}$$

$$(S_E^*, S_F^*, S_G^*) = \begin{pmatrix} I & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & I & 0 & 0 & 0 & 0 & 0 & 0 & I & 0 & n_1 \\ 0 & I & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & I & 0 & 0 & 0 & 0 & 0 & 0 & 0 & n_2 \\ 0 & 0 & I & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & I & 0 & 0 & n_3 \\ 0 & 0 & 0 & I & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & I & 0 & 0 & 0 & 0 & n_4 \\ 0 & 0 & 0 & 0 & I & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & n_5 \\ 0 & 0 & 0 & 0 & 0 & 0 & I & 0 & 0 & 0 & 0 & 0 & 0 & I & 0 & 0 & 0 & 0 & n_4 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & I & 0 & 0 & 0 & 0 & 0 & 0 & I & 0 & 0 & 0 & n_6 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & I & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & n_7 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & I & 0 & 0 & 0 & 0 & 0 & n_8 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & n - r_{e|f|g} - t \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & t \end{pmatrix}, \tag{2.5}$$

where

$$t = r_{abcd|e000|f000|g000} - r_{bcd} - r_{e|f|g},$$

$$m_1 = r_b + r_c + r_d - r_{db0|d0c},\ m_2 = r_{db0|d0c} - r_{bc} - r_d, \tag{2.6}$$

$$m_3 = r_{db0|d0c} - r_{bd} - r_c,\ m_4 = r_{bc} + r_{cd} + r_{bd} - r_{bcd} - r_{db0|d0c}, \tag{2.7}$$

$$m_5 = r_{bcd} - r_{cd},\ m_6 = r_{db0|d0c} - r_{cd} - r_b,\ m_7 = r_{bcd} - r_{bd},\ m_8 = r_{bcd} - r_{bc}, \tag{2.8}$$

$$n_1 = r_e + r_f + r_g - r_{gg|e0|0f},\ n_2 = r_{gg|e0|0f} - r_{e|f} - r_g, \tag{2.9}$$

$$n_3 = r_{gg|e0|0f} - r_{e|f} - r_f,\ n_4 = r_{e|f} + r_{f|g} + r_{e|g} - r_{e|f|g} - r_{gg|e0|0f}, \tag{2.10}$$

$$n_5 = r_{e|f|g} - r_{f|g},\ n_6 = r_{gg|e0|0f} - r_{f|g} - r_e,\ n_7 = r_{e|f|g} - r_{e|g},\ n_8 = r_{e|f|g} - r_{e|f}. \tag{2.11}$$

The block columns of S_A are $(n_1, n_2, n_3, n_4, n_5, n_4, n_6.n_7, n_8, n - r_{e|f|g} - t, t)$. The block rows of S_A are $(m_1, m_2, m_3, m_4, m_5, m_4, m_6.m_7, m_8, m - r_{bcd} - t, t)$.

Proof. The proof is constructive. We establish the result through the following steps. First, we give equivalence canonical forms of the general matrix arrays $(B,\ C,\ D)$ and $\begin{pmatrix} E \\ F \\ G \end{pmatrix}$. Second, we provide the simultaneous decomposition of the matrix array (1.1).

Step 1. For the matrix arrays $(B,\ C,\ D)$ and $\begin{pmatrix} E \\ F \\ G \end{pmatrix}$, we can find eight matrices $P_1 \in GL_m(\mathbb{H})$, $Q_1 \in GL_n(\mathbb{H})$, $W_B \in GL_{p_1}(\mathbb{H})$, $W_C \in GL_{p_2}(\mathbb{H})$, $W_D \in GL_{p_3}(\mathbb{H})$, $W_E \in GL_{q_1}(\mathbb{H})$, $W_F \in GL_{q_2}(\mathbb{H})$, $W_G \in GL_{q_3}(\mathbb{H})$ such that

$$P_1 \begin{pmatrix} B & C & D \end{pmatrix} \begin{pmatrix} W_B & 0 & 0 \\ 0 & W_C & 0 \\ 0 & 0 & W_D \end{pmatrix} =$$

$$\begin{array}{c} \begin{pmatrix} I & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & I & 0 & 0 & 0 & 0 & 0 & 0 & I & 0 \\ 0 & I & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & I & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & I & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & I & 0 & 0 \\ 0 & 0 & 0 & I & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & I & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & I & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & I & 0 & 0 & 0 & 0 & 0 & 0 & I & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & I & 0 & 0 & 0 & 0 & 0 & 0 & I & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & I & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & I & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix} \begin{matrix} m_1 \\ m_2 \\ m_3 \\ m_4 \\ m_5 \\ m_4 \\ m_6 \\ m_7 \\ m_8 \\ m - r_{bcd} \end{matrix} \end{array},$$

$$\left[\begin{pmatrix} W_E & 0 & 0 \\ 0 & W_F & 0 \\ 0 & 0 & W_G \end{pmatrix} \begin{pmatrix} E \\ F \\ G \end{pmatrix} Q_1 \right]^* =$$

$$\begin{pmatrix}
I & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & I & 0 & 0 & 0 & 0 & 0 & 0 & I & 0 \\
0 & I & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & I & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & I & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & I & 0 & 0 \\
0 & 0 & 0 & I & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & I & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & I & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & I & 0 & 0 & 0 & 0 & 0 & 0 & I & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & I & 0 & 0 & 0 & 0 & 0 & 0 & I & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & I & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & I & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0
\end{pmatrix}
\begin{matrix}
n_1 \\ n_2 \\ n_3 \\ n_4 \\ n_5 \\ n_4 \\ n_6 \\ n_7 \\ n_8 \\ n - r_{e|f|g}
\end{matrix}.$$

Let

$$P_1 A Q_1 \triangleq \begin{pmatrix} A_{11}^{(1)} & \cdots & A_{1,10}^{(1)} \\ \vdots & \ddots & \vdots \\ A_{10,1}^{(1)} & \cdots & A_{10,10}^{(1)} \end{pmatrix},$$

where the symbol $\triangleq$ means "equals by definition". For the matrix $A_{10,10}^{(1)}$, we can find $P_2 \in GL_{m-r_{bcd}}(\mathbb{H})$, $Q_2 \in GL_{n-r_{e|f|g}}(\mathbb{H})$ such that

$$P_2 A_{10,10}^{(1)} Q_2 = \begin{pmatrix} 0 & 0 \\ 0 & I_t \end{pmatrix}, t \triangleq r(A_{10,10}^{(1)}).$$

Then we have

$$\begin{pmatrix} I_{r_{bcd}} & 0 \\ 0 & P_2 \end{pmatrix} \begin{pmatrix} A_{11}^{(1)} & \cdots & A_{1,10}^{(1)} \\ \vdots & \ddots & \vdots \\ A_{10,1}^{(1)} & \cdots & A_{10,10}^{(1)} \end{pmatrix} \begin{pmatrix} I_{r_{e|f|g}} & 0 \\ 0 & Q_2 \end{pmatrix} \triangleq \begin{pmatrix} A_{11}^{(2)} & \cdots & A_{19}^{(2)} & A_{1,10}^{(2)} & A_{1,11}^{(2)} \\ \vdots & \ddots & \vdots & \vdots & \vdots \\ A_{91}^{(2)} & \cdots & A_{99}^{(2)} & A_{9,10}^{(2)} & A_{9,11}^{(2)} \\ A_{10,1}^{(2)} & \cdots & A_{10,9}^{(2)} & 0 & 0 \\ A_{11,1}^{(2)} & \cdots & A_{11,9}^{(2)} & 0 & I_t \end{pmatrix},$$

$$\begin{pmatrix} I_{r_{bcd}} & 0 \\ 0 & P_2 \end{pmatrix} P_1 \begin{pmatrix} B & C & D \end{pmatrix} \begin{pmatrix} W_B & 0 & 0 \\ 0 & W_C & 0 \\ 0 & 0 & W_D \end{pmatrix} =$$

$$\begin{pmatrix}
I & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & I & 0 & 0 & 0 & 0 & 0 & 0 & I & 0 \\
0 & I & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & I & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & I & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & I & 0 & 0 \\
0 & 0 & 0 & I & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & I & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & I & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & I & 0 & 0 & 0 & 0 & 0 & 0 & I & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & I & 0 & 0 & 0 & 0 & 0 & 0 & I & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & I & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & I & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0
\end{pmatrix}
\begin{matrix}
m_1 \\ m_2 \\ m_3 \\ m_4 \\ m_5 \\ m_4 \\ m_6 \\ m_7 \\ m_8 \\ m - r_{bcd} - t \\ t
\end{matrix},$$

$$\left[\begin{pmatrix} W_E & 0 & 0 \\ 0 & W_F & 0 \\ 0 & 0 & W_G \end{pmatrix} \begin{pmatrix} E \\ F \\ G \end{pmatrix} Q_1 \begin{pmatrix} I_{r_{e|f|g}} & 0 \\ 0 & Q_2 \end{pmatrix}\right]^* =$$

$$\begin{pmatrix}
I & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & I & 0 & 0 & 0 & 0 & 0 & 0 & I & 0 \\
0 & I & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & I & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & I & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & I & 0 & 0 \\
0 & 0 & 0 & I & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & I & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & I & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & I & 0 & 0 & 0 & 0 & 0 & 0 & I & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & I & 0 & 0 & 0 & 0 & 0 & 0 & I & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & I & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & I & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0
\end{pmatrix}
\begin{matrix}
n_1 \\ n_2 \\ n_3 \\ n_4 \\ n_5 \\ n_4 \\ n_6 \\ n_7 \\ n_8 \\ n - r_{e|f|g} - t \\ t
\end{matrix}.$$

Let

$$P_3 = \begin{pmatrix} I_{r_{bcd}} & \begin{pmatrix} 0 & -A^{(2)}_{1,11} \\ \vdots & \vdots \\ 0 & -A^{(2)}_{9,11} \end{pmatrix} \\ 0 & I_{m-r_{bcd}} \end{pmatrix},\ Q_3 = \begin{pmatrix} I_{r_{e|f|g}} & 0 \\ \begin{pmatrix} 0 & \cdots & 0 \\ -A^{(2)}_{11,1} & \cdots & -A^{(2)}_{11,9} \end{pmatrix} & I_{n-r_{e|f|g}} \end{pmatrix}.$$

Then we obtain

$$P_3 \begin{pmatrix} A^{(2)}_{11} & \cdots & A^{(2)}_{19} & A^{(2)}_{1,10} & A^{(2)}_{1,11} \\ \vdots & \ddots & \vdots & \vdots & \vdots \\ A^{(2)}_{91} & \cdots & A^{(2)}_{99} & A^{(2)}_{9,10} & A^{(2)}_{9,11} \\ A^{(2)}_{10,1} & \cdots & A^{(2)}_{10,9} & 0 & 0 \\ A^{(2)}_{11,1} & \cdots & A^{(2)}_{11,9} & 0 & I_t \end{pmatrix} Q_3 \triangleq \begin{pmatrix} A_{11} & \cdots & A_{19} & A_{1,10} & 0 \\ \vdots & \ddots & \vdots & \vdots & \vdots \\ A_{91} & \cdots & A_{99} & A_{9,10} & 0 \\ A_{10,1} & \cdots & A_{10,9} & 0 & 0 \\ 0 & \cdots & 0 & 0 & I_t \end{pmatrix}.$$

Let

$$P \triangleq P_3 \begin{pmatrix} I_{r_{bcd}} & 0 \\ 0 & P_2 \end{pmatrix} P_1,\ Q \triangleq Q_1 \begin{pmatrix} I_{r_{e|f|g}} & 0 \\ 0 & Q_2 \end{pmatrix} Q_3,$$

$$T_1 = W_C,\ T_2 = W_D,\ T_3 = W_E,\ V_1 = W_E,\ V_2 = W_F,\ V_3 = W_G.$$

Hence, the matrices $P \in GL_m(\mathbb{H})$, $Q \in GL_n(\mathbb{H})$, $T_1 \in GL_{p_1}(\mathbb{H})$, $T_2 \in GL_{p_2}(\mathbb{H})$, $T_3 \in GL_{p_3}(\mathbb{H})$, $V_1 \in GL_{q_1}(\mathbb{H})$, $V_2 \in GL_{q_2}(\mathbb{H})$, $V_3 \in GL_{q_3}(\mathbb{H})$ satisfy the equation (2.2). It follows from $S_A, S_B, S_C, S_D, S_E, S_F$, and S_G in (2.3)-(2.5) that

$$\begin{pmatrix} 1 & 1 & 1 & 1 & 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 1 & 0 & 1 & 1 & 0 \\ 1 & 0 & 1 & 1 & 0 & 1 & 0 & 1 \\ 1 & 1 & 1 & 2 & 1 & 1 & 1 & 0 \\ 1 & 1 & 1 & 2 & 1 & 1 & 0 & 1 \\ 1 & 1 & 1 & 2 & 0 & 1 & 1 & 1 \\ 1 & 1 & 1 & 2 & 1 & 1 & 1 & 1 \\ 1 & 0 & 1 & 1 & 0 & 1 & 0 & 1 \end{pmatrix} \begin{pmatrix} m_1 \\ m_2 \\ m_3 \\ m_4 \\ m_5 \\ m_6 \\ m_7 \\ m_8 \end{pmatrix} = \begin{pmatrix} r_b \\ r_c \\ r_d \\ r_{bc} \\ r_{bd} \\ r_{cd} \\ r_{bcd} \\ r_{db0|d0c} - r_b - r_c \end{pmatrix},$$

$$\begin{pmatrix} 1 & 1 & 1 & 1 & 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 1 & 0 & 1 & 1 & 0 \\ 1 & 0 & 1 & 1 & 0 & 1 & 0 & 1 \\ 1 & 1 & 1 & 2 & 1 & 1 & 1 & 0 \\ 1 & 1 & 1 & 2 & 1 & 1 & 0 & 1 \\ 1 & 1 & 1 & 2 & 0 & 1 & 1 & 1 \\ 1 & 1 & 1 & 2 & 1 & 1 & 1 & 1 \\ 1 & 0 & 1 & 1 & 0 & 1 & 0 & 1 \end{pmatrix} \begin{pmatrix} n_1 \\ n_2 \\ n_3 \\ n_4 \\ n_5 \\ n_6 \\ n_7 \\ n_8 \end{pmatrix} = \begin{pmatrix} r_e \\ r_f \\ r_g \\ r_{e|f} \\ r_{e|g} \\ r_{f|g} \\ r_{e|f|g} \\ r_{gg|e0|0f} - r_e - r_f \end{pmatrix}.$$

Solving for $m_i, n_i, (i = 1, \ldots, 8)$ gives (2.6)-(2.11). □

Remark 2.1. *Wang et. al.* [19] *did not give the values of* m_i, $(i = 1, 2, \ldots, 8)$ *in Lemma 2.1. As a special case of Theorem 2.2, we can derive all the dimensions of identity matrices in the equivalence canonical form of triple real quaternion matrices* (B, C, D)*, i.e., the values of* m_i, $(i = 1, 2, \ldots, 8)$ *in Lemma 2.1:*

$$m_1 = r_b + r_c + r_d - r_{db0|d0c},\ m_2 = r_{db0|d0c} - r_{bc} - r_d,$$

$$m_3 = r_{db0|d0c} - r_{bd} - r_c,\ m_4 = r_{bc} + r_{cd} + r_{bd} - r_{bcd} - r_{db0|d0c},$$

$$m_5 = r_{bcd} - r_{cd},\ m_6 = r_{db0|d0c} - r_{cd} - r_b,\ m_7 = r_{bcd} - r_{bd},\ m_8 = r_{bcd} - r_{bc}.$$

On the other hand, the values of $m_i, n_i, (i = 1, \ldots, 8)$ *play an important role in investigating the range of ranks of the general solutions to (1.2) and (1.3).*

3. **Some applications of the simultaneous decomposition of (1.1)**

The simultaneous decomposition of (1.1) is useful in solving the following questions:

- §3.2. Give some solvability conditions and an expression of the general solution to the real quaternion matrix equation (1.2).
- §3.3. Give the range of ranks of the general solution in the real quaternion matrix equation (1.2).
- §3.4. Give some solvability conditions and an expression of the general solution to the real quaternion matrix equation (1.3).
- §3.5. Give the range of ranks of the general solution in the real quaternion matrix equation (1.3).

3.1. **Preliminaries.** In this section, we give some lemmas which are used in the further development of this paper. The following Lemmas are due to [7], [8] and [22]-[24] which can be generalized to $\mathbb{H}$.

Lemma 3.1. *(*[7]-[9], [22]-[24]*) Let*

$$H(X,Y)=\begin{pmatrix} A_1 & B_1 & C_1 \\ D_1 & X & E_1 \\ F_1 & G_1 & Y \end{pmatrix}, \tag{3.1}$$

where $A_1\in\mathbb{H}^{\tilde{n}\times n}, B_1\in\mathbb{H}^{\tilde{n}\times m}, C_1\in\mathbb{H}^{\tilde{n}\times p}, D_1\in\mathbb{H}^{\tilde{m}\times n}, E_1\in\mathbb{H}^{\tilde{m}\times p}, F_1\in\mathbb{H}^{\tilde{p}\times n}$ *and* $G_1\in\mathbb{H}^{\tilde{p}\times m}$ *are given, and* $X\in\mathbb{H}^{\tilde{m}\times m}$ *and* $Y\in\mathbb{H}^{\tilde{p}\times p}$ *are variable matrices. Then,*

$$\max_{X,Y} r\left[H\left(X,Y\right)\right]=\min\left\{\tilde{m}+\tilde{p}+r_{a_1b_1c_1},\tilde{m}+p+r_{a_1b_1|f_1g_1},m+\tilde{p}+r_{a_1c_1|d_1e_1},m+p+r_{a_1|d_1|f_1}\right\},$$

$$\min_{X,Y} r\left[H\left(X,Y\right)\right]=r_{a_1b_1c_1}+r_{a_1|d_1|f_1}+\max\left\{r_{a_1c_1|d_1e_1}-r_{a_1c_1}-r_{a_1|d_1},r_{a_1b_1|f_1g_1}-r_{a_1b_1}-r_{a_1|f_1}\right\}.$$

Lemma 3.2. *(*[7]-[9], [22]-[24]*) Let*

$$M(X,Y)=\begin{pmatrix} A_1 & X \\ Y & B_1 \end{pmatrix} \tag{3.2}$$

where $A_1\in\mathbb{H}^{m\times n}$ *and* $B_1\in\mathbb{H}^{p\times q}$ *are given, and* $X\in\mathbb{H}^{m\times q}$ *and* $Y\in\mathbb{H}^{p\times n}$ *are variable matrices. Then,*

$$\max_{X,Y} r\left[M\left(X,Y\right)\right]=\min\left\{m+p,n+q,r_{a_1}+p+q,r_{b_1}+m+n\right\},$$

$$\min_{X,Y} r\left[M\left(X,Y\right)\right]=\max\left\{r_{a_1},\ r_{b_1}\right\}.$$

Lemma 3.3. *(*[8], [9], [22]-[24]*) Let*

$$M_2=\begin{pmatrix} Y & D_1 \\ B_1 & A_1 \end{pmatrix},$$

where A_1, B_1 *and* D_1 *are given, and* $Y\in\mathbb{H}^{n\times m}$ *is a variable matrix. Then,*

$$\max_{Y\in\mathbb{H}^{n\times m}} r\left(M_2\right)=\min\left\{n+r_{a_1b_1},m+r_{a_1|d_1}\right\},\ \min_{Y\in\mathbb{H}^{n\times m}} r\left(M_2\right)=r_{a_1b_1}+r_{a_1|d_1}-r_{a_1}.$$

3.2. **Some solvability conditions and the general solution to (1.2).** In this section, the simultaneous decomposition of (1.1) will be used to solve the real quaternion matrix equation (1.2).

Theorem 3.4. *Let $A \in \mathbb{H}^{m\times n}, B \in \mathbb{H}^{m\times p_1}, C \in \mathbb{H}^{m\times p_2}, D \in \mathbb{H}^{m\times p_3}, E \in \mathbb{H}^{q_1\times n}, F \in \mathbb{H}^{q_2\times n}$ and $G \in \mathbb{H}^{q_3\times n}$ be given. Then the equation (1.2) is consistent if and only if*

$$A_{94} = A_{96},\ A_{49} = A_{69},\ A_{64} = A_{46},$$

$$r_{abcd|e000|f000|g000} = r_{bcd} + r_{e|f|g},\quad \left(A^*_{1,10},\ \cdots,\ A^*_{9,10}\right) = 0,\quad (A_{10,1},\ \cdots,\ A_{10,9}) = 0,$$

$$A_{29} = 0,\ A_{92} = 0,\ A_{38} = 0,\ A_{83} = 0,\ A_{48} = 0,\ A_{84} = 0,\ A_{56} = 0,\ A_{65} = 0,$$

$$A_{57} = 0,\ A_{75} = 0,\ A_{58} = 0,\ A_{85} = 0,\ A_{59} = 0,\ A_{95} = 0,\ A_{89} = 0,\ A_{98} = 0. \tag{3.3}$$

In this case, the general solution to (1.2) can be expressed as

$$X = T_1^{-1}\widehat{X}V_1^{-1},\quad Y = T_2^{-1}\widehat{Y}V_2^{-1},\quad Z = T_3^{-1}\widehat{Z}V_3^{-1},$$

where

$$\widehat{X} = \begin{array}{c} \\ m_1 \\ m_2 \\ m_3 \\ m_4 \\ m_5 \\ p_1 - r_b \end{array} \begin{array}{c} \begin{array}{cccccc} n_1 & n_2 & n_3 & n_4 & n_5 & q_1 - r_e \end{array} \\ \begin{pmatrix} X_{11} & X_{12} & X_{13} & X_{14} & A_{15} & X_{16} \\ X_{21} & X_{22} & A_{23} & A_{24} & A_{25} & X_{26} \\ X_{31} & A_{32} & X_{33} & A_{34} - A_{36} & A_{35} & X_{36} \\ X_{41} & A_{42} & A_{43} - A_{63} & A_{44} - A_{64} & A_{45} & X_{46} \\ A_{51} & A_{52} & A_{53} & A_{54} & A_{55} & X_{56} \\ X_{61} & X_{62} & X_{63} & X_{64} & X_{65} & X_{66} \end{pmatrix} \end{array}, \tag{3.4}$$

$$\widehat{Y} = \begin{array}{c} \\ m_4 \\ m_6 \\ m_7 \\ m_1 \\ m_2 \\ p_2 - r_c \end{array} \begin{array}{c} \begin{array}{cccccc} n_4 & n_6 & n_7 & n_1 & n_2 & q_2 - r_f \end{array} \\ \begin{pmatrix} A_{66} - A_{64} & A_{67} - A_{47} & A_{68} & A_{61} - A_{41} + X_{41} & A_{62} & Y_{16} \\ A_{76} - A_{74} & Y_{22} & A_{78} & Y_{24} & A_{72} & Y_{26} \\ A_{86} & A_{87} & A_{88} & A_{81} & A_{82} & Y_{36} \\ A_{16} - A_{14} + X_{14} & Y_{42} & A_{18} & Y_{44} & A_{12} - X_{12} & Y_{46} \\ A_{26} & A_{27} & A_{28} & A_{21} - X_{21} & A_{22} - X_{22} & Y_{56} \\ Y_{61} & Y_{62} & Y_{63} & Y_{64} & Y_{65} & Y_{66} \end{pmatrix} \end{array}, \tag{3.5}$$

$$\widehat{Z} = \begin{array}{c} \\ m_8 \\ m_4 \\ m_6 \\ m_3 \\ m_1 \\ p_3 - r_d \end{array} \begin{array}{c} \begin{array}{cccccc} n_8 & n_4 & n_6 & n_3 & n_1 & q_3 - r_g \end{array} \\ \begin{pmatrix} A_{99} & A_{96} & A_{97} & A_{93} & A_{91} & Z_{16} \\ A_{69} & A_{64} & A_{47} & A_{63} & A_{41} - X_{41} & Z_{26} \\ A_{79} & A_{74} & A_{77} - Y_{22} & A_{73} & A_{71} - Y_{24} & Z_{36} \\ A_{39} & A_{36} & A_{37} & A_{33} - X_{33} & A_{31} - X_{31} & Z_{46} \\ A_{19} & A_{14} - X_{14} & A_{17} - Y_{42} & A_{13} - X_{13} & Z_{55} & Z_{56} \\ Z_{61} & Z_{62} & Z_{63} & Z_{64} & Z_{65} & Z_{66} \end{pmatrix} \end{array}, \tag{3.6}$$

A_{ij}, T_i, V_i are defined in Theorem 2.2, the remaining X_{ij}, Y_{ij}, Z_{ij} in (3.4)-(3.6) are arbitrary matrices over $\mathbb{H}$ with appropriate sizes.

Proof. It follows from Theorem 2.2 that the matrix equation (1.2) is equivalent to the matrix equation

$$S_B(T_1XV_1)S_E + S_C(T_2YV_2)S_F + S_D(T_3ZV_3)S_G = S_A. \tag{3.7}$$

Let the matrices

$$\widehat{X} = T_1XV_1 = \begin{pmatrix} X_{11} & \cdots & X_{16} \\ \vdots & \ddots & \vdots \\ X_{61} & \cdots & X_{66} \end{pmatrix}, \tag{3.8}$$

$$\widehat{Y} = T_2YV_2 = \begin{pmatrix} Y_{11} & \cdots & Y_{16} \\ \vdots & \ddots & \vdots \\ Y_{61} & \cdots & Y_{66} \end{pmatrix}, \tag{3.9}$$

$$\widehat{Z} = T_3ZV_3 = \begin{pmatrix} Z_{11} & \cdots & Z_{16} \\ \vdots & \ddots & \vdots \\ Z_{61} & \cdots & Z_{66} \end{pmatrix}, \tag{3.10}$$

be partitioned in accordance with (3.7). Then it follows from (2.3)-(2.5) and (3.7)-(3.10) that

$$\begin{pmatrix} X_{11}+Y_{44}+Z_{55} & X_{12}+Y_{45} & X_{13}+Z_{54} & X_{14}+Z_{52} & X_{15} & Y_{41}+Z_{52} & Y_{42}+Z_{53} & Y_{43} & Z_{51} & 0 & 0 \\ X_{21}+Y_{54} & X_{22}+Y_{55} & X_{23} & X_{24} & X_{25} & Y_{51} & Y_{52} & Y_{53} & 0 & 0 & 0 \\ X_{31}+Z_{45} & X_{32} & X_{33}+Z_{44} & X_{34}+Z_{42} & X_{35} & Z_{42} & Z_{43} & 0 & Z_{41} & 0 & 0 \\ X_{41}+Z_{25} & X_{42} & X_{43}+Z_{24} & X_{44}+Z_{22} & X_{45} & Z_{22} & Z_{23} & 0 & Z_{21} & 0 & 0 \\ X_{51} & X_{52} & X_{53} & X_{54} & X_{55} & 0 & 0 & 0 & 0 & 0 & 0 \\ Y_{14}+Z_{25} & Y_{15} & Z_{24} & Z_{22} & 0 & Y_{11}+Z_{22} & Y_{12}+Z_{23} & Y_{13} & Z_{21} & 0 & 0 \\ Y_{24}+Z_{35} & Y_{25} & Z_{34} & Z_{32} & 0 & Y_{21}+Z_{32} & Y_{22}+Z_{33} & Y_{23} & Z_{31} & 0 & 0 \\ Y_{34} & Y_{35} & 0 & 0 & 0 & Y_{31} & Y_{32} & Y_{33} & 0 & 0 & 0 \\ Z_{15} & 0 & Z_{14} & Z_{12} & 0 & Z_{12} & Z_{13} & 0 & Z_{11} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

$$= \begin{pmatrix} A_{11} & A_{12} & A_{13} & A_{14} & A_{15} & A_{16} & A_{17} & A_{18} & A_{19} & A_{1,10} & 0 \\ A_{21} & A_{22} & A_{23} & A_{24} & A_{25} & A_{26} & A_{27} & A_{28} & A_{29} & A_{2,10} & 0 \\ A_{31} & A_{32} & A_{33} & A_{34} & A_{35} & A_{36} & A_{37} & A_{38} & A_{39} & A_{3,10} & 0 \\ A_{41} & A_{42} & A_{43} & A_{44} & A_{45} & A_{46} & A_{47} & A_{48} & A_{49} & A_{4,10} & 0 \\ A_{51} & A_{52} & A_{53} & A_{54} & A_{55} & A_{56} & A_{57} & A_{58} & A_{59} & A_{5,10} & 0 \\ A_{61} & A_{62} & A_{63} & A_{64} & A_{65} & A_{66} & A_{67} & A_{68} & A_{69} & A_{6,10} & 0 \\ A_{71} & A_{72} & A_{73} & A_{74} & A_{75} & A_{76} & A_{77} & A_{78} & A_{79} & A_{7,10} & 0 \\ A_{81} & A_{82} & A_{83} & A_{84} & A_{85} & A_{86} & A_{87} & A_{88} & A_{89} & A_{8,10} & 0 \\ A_{91} & A_{92} & A_{93} & A_{94} & A_{95} & A_{96} & A_{97} & A_{98} & A_{99} & A_{9,10} & 0 \\ A_{10,1} & A_{10,2} & A_{10,3} & A_{10,4} & A_{10,5} & A_{10,6} & A_{10,7} & A_{10,8} & A_{10,9} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & I_t \end{pmatrix}. \tag{3.11}$$

If the equation (1.2) has a solution (X, Y, Z), then by (3.11), we have that the equalities in (3.3) hold, and

$X_{11} + Y_{44} + Z_{55} = A_{11},\ X_{12} + Y_{45} = A_{12},\ X_{13} + Z_{54} = A_{13},\ X_{14} + Z_{52} = A_{14},\ X_{15} = A_{15},$
$Y_{41} + Z_{52} = A_{16},\ Y_{42} + Z_{53} = A_{17},\ Y_{43} = A_{18},\ Z_{51} = A_{19}, X_{21} + Y_{54} = A_{21},\ X_{22} + Y_{55} = A_{22},$
$X_{23} = A_{23},\ X_{24} = A_{24},\ X_{25} = A_{25},\ Y_{51} = A_{26},\ Y_{52} = A_{27},\ Y_{53} = A_{28},\ X_{31} + Z_{45} = A_{31},$
$X_{32} = A_{32},\ X_{33} + Z_{44} = A_{33},\ X_{34} + Z_{42} = A_{34},\ X_{35} = A_{35},\ Z_{42} = A_{36},\ Z_{43} = A_{37},\ Z_{41} = A_{39},$
$X_{41} + Z_{25} = A_{41},\ X_{42} = A_{42},\ X_{43} + Z_{24} = A_{43},\ X_{44} + Z_{22} = A_{44},\ X_{45} = A_{45},\ Z_{22} = A_{46},$
$Z_{23} = A_{47},\ Z_{21} = A_{49},\ X_{51} = A_{51},\ X_{52} = A_{52},\ X_{53} = A_{53},\ X_{54} = A_{54},\ X_{55} = A_{55}$
$Y_{14} + Z_{25} = A_{61},\ Y_{15} = A_{62},\ Z_{24} = A_{63},\ Z_{22} = A_{64},\ Y_{11} + Z_{22} = A_{66},\ Y_{12} + Z_{23} = A_{67},$
$Y_{13} = A_{68},\ Z_{21} = A_{69},\ Y_{24} + Z_{35} = A_{71},\ Y_{25} = A_{72},\ Z_{34} = A_{73},\ Z_{32} = A_{74},\ Y_{21} + Z_{32} = A_{76},$
$Y_{22} + Z_{33} = A_{77},\ Y_{23} = A_{78},\ Z_{31} = A_{79},\ Y_{34} = A_{81},\ Y_{35} = A_{82},\ Y_{31} = A_{86},\ Y_{32} = A_{87},$
$Y_{33} = A_{88},\ Z_{15} = A_{91},\ Z_{14} = A_{93},\ Z_{12} = A_{94},\ Z_{12} = A_{96},\ Z_{13} = A_{97},\ Z_{11} = A_{99}.$

Hence, (X, Y, Z) can be expressed as (3.4)-(3.6) by (3.8)-(3.10).

Conversely, assume that the equalities in (3.3) hold, then by (2.3)-(2.5) and (3.7)-(3.11), it can be verified that the matrices have the forms of (3.4)-(3.6) is a solution of (3.7), i.e., (1.2).

□

Remark 3.1. *In our opinion the presented expression of the general solution is more useful than the expression found by Wang et al.* [19]*, since the latter can not be used to consider the maximal and minimal ranks of the general solution to (1.2).*

3.3. **The range of ranks of the general solution to (1.2).** In this section, we consider the maximal and minimal ranks of the general solution to the real quaternion matrix equation (1.2).

Theorem 3.5. *Let $A \in \mathbb{H}^{m\times n}, B \in \mathbb{H}^{m\times p_1}, C \in \mathbb{H}^{m\times p_2}, D \in \mathbb{H}^{m\times p_3}, E \in \mathbb{H}^{q_1\times n}, F \in \mathbb{H}^{q_2\times n}$ and $G \in \mathbb{H}^{q_3\times n}$ be given. Assume that equation (1.2) is consistent. Then,*

$$\max_{BXE+CYF+DZG=A} r(Z) = \min\left\{p_3,\ q_3,\ p_3+q_3+r_{a|e|f}-r_d-r_{e|f|g},\ p_3+q_3+r_{abc}-r_g-r_{bcd},\right.$$

$$\left. p_3+q_3+r_{d0ab0|da00c|0e000|00f00}-r_{gg|e0|0f}-r_{bd}-r_{cd},\ p_3+q_3+r_{ac|e0}-r_{cd}-r_{e|f},\right.$$

$$p_3+q_3+r_{ab|f0}-r_{f|g}-r_{bd},\ p_3+q_3+r_{gg00|0ab0|a00c|e000|0f00}-r_{db0|d0c}-r_{e|g}-r_{f|g}\big\},$$

$$\begin{aligned}
&\min_{BXE+CYF+DZG=A} r\left(Z\right)\\
=&r_{d0ab0|da00c|0e000|00f00}+r_{gg00|0ab0|a00c|e000|0f00}+r_{abc}+r_{a|e|f}+r_{bd}+r_{e|g}+r_{bc}+r_{e|f}\\
&-r_{d0ab00|da00cb|0e0000|00f000}-r_{gg00|0ab0|a00c|e000|0f00|0e00}\\
&+\max\big\{r_{ac|e0}-r_{d0ab0|da00c|0e000|00f00|00e00}-r_{gg000|0ab0c|a00c0|e0000|0f000},\\
&\qquad r_{ab|f0}-r_{d0ab0|da00c|0e000|00f00|0f000}-r_{gg000|0ab00|a00cb|e0000|0f000}\big\}.
\end{aligned}$$

Proof. It follows from Theorem 3.4 that the expression of Z in (1.2) can be expressed as $Z=T_3^{-1}\widehat{Z}V_3^{-1}$, where $\widehat{Z}$ is given in (3.6). Clearly, $r(Z)=r(T_3^{-1}\widehat{Z}V_3^{-1})=r(\widehat{Z})$. Now we consider the maximal and minimal ranks of $\widehat{Z}$. Applying Lemma 3.2 to the variable matrices $(Z_{61},\ Z_{62},\ Z_{63},\ Z_{64},\ Z_{65})$ and $\begin{pmatrix}Z_{16}\\Z_{26}\\Z_{36}\\Z_{46}\\Z_{56}\end{pmatrix}$ of $\widehat{Z}$, we obtain

$$\max_{Z_{i6},\ Z_{6i},\ i=1,\ldots,5} r(\widehat{Z})=\min\left\{p_3,q_3,p_3+q_3-r_d-r_g+r(\Theta),r(Z_{66})+r_d+r_g\right\},$$

$$\min_{Z_{i6},\ Z_{6i},\ i=1,\ldots,5} r(\widehat{Z})=\max\left\{r(\Theta),r(Z_{66})\right\},$$

where

$$\Theta=\begin{array}{c}\\m_8\\m_4\\m_6\\m_3\\m_1\end{array}\begin{array}{c}\begin{array}{ccccc}n_8&n_4&n_6&n_3&n_1\end{array}\\\begin{pmatrix}A_{99}&A_{96}&A_{97}&A_{93}&A_{91}\\A_{69}&A_{64}&A_{47}&A_{63}&A_{41}-X_{41}\\A_{79}&A_{74}&A_{77}-Y_{22}&A_{73}&A_{71}-Y_{24}\\A_{39}&A_{36}&A_{37}&A_{33}-X_{33}&A_{31}-X_{31}\\A_{19}&A_{14}-X_{14}&A_{17}-Y_{42}&A_{13}-X_{13}&Z_{55}\end{pmatrix}\end{array}.$$

Note that

$$\max r\left(Z_{66}\right)=\min\left\{p_3-r_d,q_3-r_g\right\},\ \max r\left(Z_{66}\right)=0.$$

Hence, we have

$$\max_{Z_{i6},\ Z_{6i},\ i=1,\ldots,6} r(\widehat{Z})=\min\left\{p_3,q_3,p_3+q_3-r_d-r_g+r(\Theta)\right\},\ \min_{Z_{i6},\ Z_{6i},\ i=1,\ldots,6} r(\widehat{Z})=r(\Theta). \tag{3.12}$$

Applying Lemma 3.3 to the variable matrices $\begin{pmatrix}A_{14}-X_{14},&A_{17}-Y_{42},&A_{13}-X_{13},&Z_{55}\end{pmatrix}$ of Θ, we obtain

$$\max_{\left(\begin{smallmatrix}A_{14}-X_{14},\ A_{17}-Y_{42},\ A_{13}-X_{13},\ Z_{55}\end{smallmatrix}\right)} r\left(\Theta\right)=\min\left\{m_1+r(\Theta_1),n_1+n_3+n_4+n_6+r\begin{pmatrix}A_{99}\\A_{69}\\A_{79}\\A_{39}\\A_{19}\end{pmatrix}\right\},$$

$$\min_{\left(\begin{smallmatrix} A_{14}-X_{14},\ A_{17}-Y_{42},\ A_{13}-X_{13},\ Z_{55} \end{smallmatrix}\right)} r\left(\Theta\right) = r(\Theta_1) + r\begin{pmatrix} A_{99} \\ A_{69} \\ A_{79} \\ A_{39} \\ A_{19} \end{pmatrix} - r\begin{pmatrix} A_{99} \\ A_{69} \\ A_{79} \\ A_{39} \end{pmatrix},$$

where

$$\Theta_1 = \begin{array}{c} \\ m_8 \\ m_4 \\ m_6 \\ m_3 \end{array} \begin{array}{c} \begin{array}{ccccc} n_8 & n_4 & n_6 & n_3 & n_1 \end{array} \\ \begin{pmatrix} A_{99} & A_{96} & A_{97} & A_{93} & A_{91} \\ A_{69} & A_{64} & A_{47} & A_{63} & A_{41}-X_{41} \\ A_{79} & A_{74} & A_{77}-Y_{22} & A_{73} & A_{71}-Y_{24} \\ A_{39} & A_{36} & A_{37} & A_{33}-X_{33} & A_{31}-X_{31} \end{pmatrix} \end{array}.$$

Applying Lemma 3.3 to the variable matrices $\begin{pmatrix} A_{41}-X_{41} \\ A_{71}-Y_{24} \\ A_{31}-X_{31} \end{pmatrix}$ of Θ_1, we obtain

$$\max_{\begin{pmatrix} A_{41}-X_{41} \\ A_{71}-Y_{24} \\ A_{31}-X_{31} \end{pmatrix}} r\left(\Theta_1\right) = \min\left\{m_3+m_4+m_6+r\begin{pmatrix} A_{99}, & A_{96}, & A_{97}, & A_{93}, & A_{91} \end{pmatrix}, n_1+r(\Theta_2)\right\},$$

$$\min_{\begin{pmatrix} A_{41}-X_{41} \\ A_{71}-Y_{24} \\ A_{31}-X_{31} \end{pmatrix}} r\left(\Theta_1\right) = r(\Theta_2) + r\begin{pmatrix} A_{99}, & A_{96}, & A_{97}, & A_{93}, & A_{91} \end{pmatrix} - r\begin{pmatrix} A_{99}, & A_{96}, & A_{97}, & A_{93} \end{pmatrix},$$

where

$$\Theta_1 = \begin{array}{c} \\ m_8 \\ m_4 \\ m_6 \\ m_3 \end{array} \begin{array}{c} \begin{array}{cccc} n_8 & n_4 & n_6 & n_3 \end{array} \\ \begin{pmatrix} A_{99} & A_{96} & A_{97} & A_{93} \\ A_{69} & A_{64} & A_{47} & A_{63} \\ A_{79} & A_{74} & A_{77}-Y_{22} & A_{73} \\ A_{39} & A_{36} & A_{37} & A_{33}-X_{33} \end{pmatrix} \end{array}.$$

Applying Lemma 3.1 to the variable matrices $A_{77}-Y_{22}$ and $A_{33}-X_{33}$ of Θ_2, we obtain

$$\begin{aligned} &\max_{A_{77}-Y_{22},\ A_{33}-X_{33}} r\left(\Theta_2\right) \\ =&\min\left\{m_3+m_6+r\begin{pmatrix} A_{99} & A_{96} & A_{97} & A_{93} \\ A_{69} & A_{64} & A_{47} & A_{63} \end{pmatrix}, n_3+m_6+r\begin{pmatrix} A_{99} & A_{96} & A_{97} \\ A_{69} & A_{64} & A_{47} \\ A_{39} & A_{36} & A_{37} \end{pmatrix},\right. \\ &\left. m_3+n_6+r\begin{pmatrix} A_{99} & A_{96} & A_{93} \\ A_{69} & A_{64} & A_{63} \\ A_{79} & A_{74} & A_{73} \end{pmatrix}, n_3+n_6+r\begin{pmatrix} A_{99} & A_{96} \\ A_{69} & A_{64} \\ A_{79} & A_{74} \\ A_{39} & A_{36} \end{pmatrix}\right\}. \end{aligned} \tag{3.13}$$

$$\min_{A_{77}-Y_{22},\ A_{33}-X_{33}} r\left(\Theta_2\right) = r\begin{pmatrix} A_{99} & A_{96} & A_{97} & A_{93} \\ A_{69} & A_{64} & A_{47} & A_{63} \end{pmatrix} + r\begin{pmatrix} A_{99} & A_{96} \\ A_{69} & A_{64} \\ A_{79} & A_{74} \\ A_{39} & A_{36} \end{pmatrix}$$

$$+ \max\left\{ r\begin{pmatrix} A_{99} & A_{96} & A_{97} \\ A_{69} & A_{64} & A_{47} \\ A_{39} & A_{36} & A_{37} \end{pmatrix} - r\begin{pmatrix} A_{99} & A_{96} & A_{97} \\ A_{69} & A_{64} & A_{47} \end{pmatrix} - r\begin{pmatrix} A_{99} & A_{96} \\ A_{69} & A_{64} \\ A_{39} & A_{36} \end{pmatrix},\right.$$

$$\left. r\begin{pmatrix} A_{99} & A_{96} & A_{93} \\ A_{69} & A_{64} & A_{63} \\ A_{79} & A_{74} & A_{73} \end{pmatrix} - r\begin{pmatrix} A_{99} & A_{96} & A_{93} \\ A_{69} & A_{64} & A_{63} \end{pmatrix} - r\begin{pmatrix} A_{99} & A_{96} \\ A_{69} & A_{64} \\ A_{79} & A_{74} \end{pmatrix} \right\}.$$

Hence,

$$\max_{BXE+CYF+DZG=A} r\left(Z\right) = \min\left\{p_3, q_3, s_1, s_2, s_3, s_4, s_5, s_6\right\},$$

$$\min_{BXE+CYF+DZG=A} r\left(Z\right) = \max\left\{s_7, s_8\right\},$$

where

$$s_1 = p_3 + q_3 - r_d - r_g + n_1 + n_3 + n_4 + n_6 + r\begin{pmatrix} A_{99} \\ A_{69} \\ A_{79} \\ A_{39} \\ A_{19} \end{pmatrix},$$

$$s_2 = p_3 + q_3 - r_d - r_g + m_1 + m_3 + m_4 + m_6 + r\left(A_{99}, \quad A_{96}, \quad A_{97}, \quad A_{93}, \quad A_{91}\right),$$

$$s_3 = p_3 + q_3 - r_d - r_g + m_1 + n_1 + m_3 + m_6 + r\begin{pmatrix} A_{99} & A_{96} & A_{97} & A_{93} \\ A_{69} & A_{64} & A_{47} & A_{63} \end{pmatrix},$$

$$s_4 = p_3 + q_3 - r_d - r_g + m_1 + n_1 + n_3 + m_6 + r\begin{pmatrix} A_{99} & A_{96} & A_{97} \\ A_{69} & A_{64} & A_{47} \\ A_{39} & A_{36} & A_{37} \end{pmatrix},$$

$$s_5 = p_3 + q_3 - r_d - r_g + m_1 + n_1 + m_3 + n_6 + r\begin{pmatrix} A_{99} & A_{96} & A_{93} \\ A_{69} & A_{64} & A_{63} \\ A_{79} & A_{74} & A_{73} \end{pmatrix},$$

$$s_6 = p_3 + q_3 - r_d - r_g + m_1 + n_1 + n_3 + n_6 + r\begin{pmatrix} A_{99} & A_{96} \\ A_{69} & A_{64} \\ A_{79} & A_{74} \\ A_{39} & A_{36} \end{pmatrix},$$

$$
\begin{aligned}
s_7 =& r\begin{pmatrix} A_{99} & A_{96} & A_{97} & A_{93} \\ A_{69} & A_{64} & A_{47} & A_{63} \end{pmatrix} + r\begin{pmatrix} A_{99} & A_{96} \\ A_{69} & A_{64} \\ A_{79} & A_{74} \\ A_{39} & A_{36} \end{pmatrix} + r\begin{pmatrix} A_{99} \\ A_{69} \\ A_{79} \\ A_{39} \\ A_{19} \end{pmatrix} - r\begin{pmatrix} A_{99} \\ A_{69} \\ A_{79} \\ A_{39} \end{pmatrix} \\
&+ r\begin{pmatrix} A_{99}, & A_{96}, & A_{97}, & A_{93}, & A_{91} \end{pmatrix} - r\begin{pmatrix} A_{99}, & A_{96}, & A_{97}, & A_{93} \end{pmatrix} \\
&+ r\begin{pmatrix} A_{99} & A_{96} & A_{97} \\ A_{69} & A_{64} & A_{47} \\ A_{39} & A_{36} & A_{37} \end{pmatrix} - r\begin{pmatrix} A_{99} & A_{96} & A_{97} \\ A_{69} & A_{64} & A_{47} \end{pmatrix} - r\begin{pmatrix} A_{99} & A_{96} \\ A_{69} & A_{64} \\ A_{39} & A_{36} \end{pmatrix},
\end{aligned}
$$

$$
\begin{aligned}
s_8 =& r\begin{pmatrix} A_{99} & A_{96} & A_{97} & A_{93} \\ A_{69} & A_{64} & A_{47} & A_{63} \end{pmatrix} + r\begin{pmatrix} A_{99} & A_{96} \\ A_{69} & A_{64} \\ A_{79} & A_{74} \\ A_{39} & A_{36} \end{pmatrix} + r\begin{pmatrix} A_{99} \\ A_{69} \\ A_{79} \\ A_{39} \\ A_{19} \end{pmatrix} - r\begin{pmatrix} A_{99} \\ A_{69} \\ A_{79} \\ A_{39} \end{pmatrix} \\
&+ r\begin{pmatrix} A_{99}, & A_{96}, & A_{97}, & A_{93}, & A_{91} \end{pmatrix} - r\begin{pmatrix} A_{99}, & A_{96}, & A_{97}, & A_{93} \end{pmatrix} \\
&+ r\begin{pmatrix} A_{99} & A_{96} & A_{93} \\ A_{69} & A_{64} & A_{63} \\ A_{79} & A_{74} & A_{73} \end{pmatrix} - r\begin{pmatrix} A_{99} & A_{96} & A_{93} \\ A_{69} & A_{64} & A_{63} \end{pmatrix} - r\begin{pmatrix} A_{99} & A_{96} \\ A_{69} & A_{64} \\ A_{79} & A_{74} \end{pmatrix}.
\end{aligned}
$$

Now we pay attention to the ranks of the block matrices in s_i. Upon construction and computation, we obtain

$$
\begin{aligned}
r\begin{pmatrix} A_{99} & A_{96} \\ A_{69} & A_{64} \\ A_{79} & A_{74} \\ A_{39} & A_{36} \end{pmatrix} =& r\begin{pmatrix} S_G & S_G & 0 & 0 \\ 0 & S_A & S_B & 0 \\ S_A & 0 & 0 & S_C \\ S_E & 0 & 0 & 0 \\ 0 & S_F & 0 & 0 \end{pmatrix} - r(S_E) - r(S_F) - r(S_B) - r(S_C) \\
&- n_3 - n_4 - n_6 - n_8 \\
=& r_{gg00|0ab0|a00c|e000|0f00} - r_e - r_f - r_b - r_c - n_3 - n_4 - n_6 - n_8, \qquad (3.14)
\end{aligned}
$$

$$
\begin{aligned}
r\begin{pmatrix} A_{99} & A_{96} & A_{97} & A_{93} \\ A_{69} & A_{64} & A_{47} & A_{63} \end{pmatrix} =& r\begin{pmatrix} S_D & 0 & S_A & S_B & 0 \\ S_D & S_A & 0 & 0 & S_C \\ 0 & S_E & 0 & 0 & 0 \\ 0 & 0 & S_F & 0 & 0 \end{pmatrix} - r(S_E) - r(S_F) - r(S_B) - r(S_C) \\
&- m_3 - m_4 - m_6 - m_8 \\
=& r_{d0ab0|da00c|0e000|00f00} - r_e - r_f - r_b - r_c - m_3 - m_4 - m_6 - m_8, \qquad (3.15)
\end{aligned}
$$

$$
r(A_{99},\ A_{96},\ A_{97},\ A_{93},\ A_{91}) = r(S_A,\ S_B,\ S_C) - r(S_B,\ S_C) = r_{abc} - r_{bc}, \qquad (3.16)
$$

$$r\begin{pmatrix}A_{99}\\A_{69}\\A_{79}\\A_{39}\\A_{19}\end{pmatrix}=r\begin{pmatrix}S_A\\S_E\\S_F\end{pmatrix}-r\begin{pmatrix}S_E\\S_F\end{pmatrix}=r_{a|e|f}-r_{e|f}, \tag{3.17}$$

$$\begin{aligned}r\begin{pmatrix}A_{99}\\A_{69}\\A_{79}\\A_{39}\end{pmatrix}=&r\begin{pmatrix}S_G&S_G&0&0\\0&S_A&S_B&0\\S_A&0&0&S_C\\S_E&0&0&0\\0&S_F&0&0\\0&S_E&0&0\end{pmatrix}-r\begin{pmatrix}S_E\\S_F\end{pmatrix}-r(S_E)-r(S_B)-r(S_C)\\&-n_4-n_6-n_8\\=&r_{gg00|0ab0|a00c|e000|0f00|0e00}-r_{e|f}-r_e-r_b-r_c-n_4-n_6-n_8,\end{aligned} \tag{3.18}$$

$$\begin{aligned}r\begin{pmatrix}A_{99},&A_{96},&A_{97},&A_{93}\end{pmatrix}=&r\begin{pmatrix}S_D&0&S_A&S_B&0&0\\S_D&S_A&0&0&S_C&S_B\\0&S_E&0&0&0&0\\0&0&S_F&0&0&0\end{pmatrix}-r(S_E)-r(S_F)-r(S_B)\\&-r(S_B,\ S_C)-m_4-m_6-m_8\\=&r_{d0ab00|da00cb|0e0000|00f000}-r_e-r_f-r_b-r_{bc}-m_4-m_6-m_8,\end{aligned} \tag{3.19}$$

$$r\begin{pmatrix}A_{99}&A_{96}&A_{93}\\A_{69}&A_{64}&A_{63}\\A_{79}&A_{74}&A_{73}\end{pmatrix}=r\begin{pmatrix}S_A&S_B\\S_F&0\end{pmatrix}-r(S_B)-r(S_F)=r_{ab|f0}-r_b-r_f, \tag{3.20}$$

$$r\begin{pmatrix}A_{99}&A_{96}&A_{97}\\A_{69}&A_{64}&A_{47}\\A_{39}&A_{36}&A_{37}\end{pmatrix}=r\begin{pmatrix}S_A&S_C\\S_E&0\end{pmatrix}-r(S_E)-r(S_C)=r_{ac|e0}-r_e-r_c, \tag{3.21}$$

$$\begin{aligned}r\begin{pmatrix}A_{99}&A_{96}\\A_{69}&A_{64}\\A_{39}&A_{36}\end{pmatrix}=&r\begin{pmatrix}S_G&S_G&0&0&0\\0&S_A&S_B&0&S_C\\S_A&0&0&S_C&0\\S_E&0&0&0&0\\0&S_F&0&0&0\end{pmatrix}-r(S_E)-r(S_F)-r(S_C)-r(S_B,\ S_C)\\&-n_3-n_4-n_6-n_8\\=&r_{gg000|0ab0c|a00c0|e0000|0f000}-r_e-r_f-r_c-r_{bc}-n_3-n_4-n_6-n_8,\end{aligned} \tag{3.22}$$

$$r\begin{pmatrix} A_{99} & A_{96} \\ A_{69} & A_{64} \\ A_{79} & A_{74} \end{pmatrix} = r\begin{pmatrix} S_G & S_G & 0 & 0 & 0 \\ 0 & S_A & S_B & 0 & 0 \\ S_A & 0 & 0 & S_C & S_B \\ S_E & 0 & 0 & 0 & 0 \\ 0 & S_F & 0 & 0 & 0 \end{pmatrix} - r(S_E) - r(S_F) - r(S_B) - r(S_B,\ S_C)$$
$$- n_3 - n_4 - n_6 - n_8$$
$$= r_{gg000|0ab00|a00cb|e0000|0f000} - r_e - r_f - r_b - r_{bc} - n_3 - n_4 - n_6 - n_8, \quad (3.23)$$

$$r\begin{pmatrix} A_{99} & A_{96} & A_{97} \\ A_{69} & A_{64} & A_{47} \end{pmatrix} = r\begin{pmatrix} S_D & 0 & S_A & S_B & 0 \\ S_D & S_A & 0 & 0 & S_C \\ 0 & S_E & 0 & 0 & 0 \\ 0 & 0 & S_F & 0 & 0 \\ 0 & 0 & S_E & 0 & 0 \end{pmatrix} - r(S_B) - r(S_C) - r(S_E) - r\begin{pmatrix} S_E \\ S_F \end{pmatrix}$$
$$- m_3 - m_4 - m_6 - m_8$$
$$= r_{d0ab0|da00c|0e000|00f00|00e00} - r_b - r_c - r_e - r_{e|f} - m_3 - m_4 - m_6 - m_8, \quad (3.24)$$

$$r\begin{pmatrix} A_{99} & A_{96} & A_{93} \\ A_{69} & A_{64} & A_{63} \end{pmatrix} = r\begin{pmatrix} S_D & 0 & S_A & S_B & 0 \\ S_D & S_A & 0 & 0 & S_C \\ 0 & S_E & 0 & 0 & 0 \\ 0 & 0 & S_F & 0 & 0 \\ 0 & S_F & 0 & 0 & 0 \end{pmatrix} - r(S_B) - r(S_C) - r(S_F) - r\begin{pmatrix} S_E \\ S_F \end{pmatrix}$$
$$- m_3 - m_4 - m_6 - m_8$$
$$= r_{d0ab0|da00c|0e000|00f00|0f000} - r_b - r_c - r_f - r_{e|f} - m_3 - m_4 - m_6 - m_8. \quad (3.25)$$

Hence from (2.6)-(2.11) and (3.14)-(3.25), we deduce that

$$s_1 = p_3 + q_3 - r_d - r_g + n_1 + n_3 + n_4 + n_6 + r\begin{pmatrix} A_{99} \\ A_{69} \\ A_{79} \\ A_{39} \\ A_{19} \end{pmatrix} = p_3 + q_3 + r_{a|e|f} - r_d - r_{e|f|g},$$

$$s_2 = p_3 + q_3 - r_d - r_g + m_1 + m_3 + m_4 + m_6 + r\left(A_{99},\ \ A_{96},\ \ A_{97},\ \ A_{93},\ \ A_{91}\right)$$
$$= p_3 + q_3 + r_{abc} - r_g - r_{bcd},$$

$$s_3 = p_3 + q_3 - r_d - r_g + m_1 + n_1 + m_3 + m_6 + r\begin{pmatrix} A_{99} & A_{96} & A_{97} & A_{93} \\ A_{69} & A_{64} & A_{47} & A_{63} \end{pmatrix}$$
$$= p_3 + q_3 - r_b - r_c - r_d - r_e - r_f - r_g + r_{d0ab0|da00c|0e000|00f00} + n_1 + m_1 - m_4 - m_8$$
$$= p_3 + q_3 + r_{d0ab0|da00c|0e000|00f00} - r_{gg|e|f} - r_{bd} - r_{cd},$$

$$
\begin{aligned}
s_4 =& p_3+q_3-r_d-r_g+m_1+n_1+n_3+m_6+r\begin{pmatrix}A_{99}&A_{96}&A_{97}\\A_{69}&A_{64}&A_{47}\\A_{39}&A_{36}&A_{37}\end{pmatrix}\\
=& p_3+q_3-r_d-r_g+m_1+n_1+n_3+m_6+r_{ac|e0}-r_e-r_c=p_3+q_3+r_{ac|e0}-r_{cd}-r_{e|f},
\end{aligned}
$$

$$
\begin{aligned}
s_5 =& p_3+q_3-r_d-r_g+m_1+n_1+m_3+n_6+r\begin{pmatrix}A_{99}&A_{96}&A_{93}\\A_{69}&A_{64}&A_{63}\\A_{79}&A_{74}&A_{73}\end{pmatrix}\\
=& p_3+q_3-r_d-r_g+m_1+n_1+m_3+n_6+r_{ab|f0}-r_b-r_f=p_3+q_3+r_{ab|f0}-r_{f|g}-r_{bd},
\end{aligned}
$$

$$
\begin{aligned}
s_6 =& p_3+q_3-r_d-r_g+m_1+n_1+n_3+n_6+r\begin{pmatrix}A_{99}&A_{96}\\A_{69}&A_{64}\\A_{79}&A_{74}\\A_{39}&A_{36}\end{pmatrix}\\
=& p_3+q_3-r_b-r_c-r_d-r_e-r_f-r_g+r_{gg00|0ab0|a00c|e000|0f00}+m_1+n_1-n_4-n_8\\
=& p_3+q_3+r_{gg00|0ab0|a00c|e000|0f00}-r_{db0|d0c}-r_{e|g}-r_{f|g},
\end{aligned}
$$

$$
\begin{aligned}
s_7 =& r_{d0ab0|da00c|0e000|00f00}+r_{gg00|0ab0|a00c|e000|0f00}+r_{abc}+r_{a|e|f}+r_{bd}+r_{e|g}+r_{bc}+r_{e|f}\\
&-r_{d0ab00|da00cb|0e0000|00f000}-r_{gg00|0ab0|a00c|e000|0f00|0e00}\\
&+r_{ac|e0}-r_{d0ab0|da00c|0e000|00f00|00e00}-r_{gg000|0ab0c|a00c0|e0000|0f000},
\end{aligned}
$$

$$
\begin{aligned}
s_8 =& r_{d0ab0|da00c|0e000|00f00}+r_{gg00|0ab0|a00c|e000|0f00}+r_{abc}+r_{a|e|f}+r_{bd}+r_{e|g}+r_{bc}+r_{e|f}\\
&-r_{d0ab00|da00cb|0e0000|00f000}-r_{gg00|0ab0|a00c|e000|0f00|0e00}\\
&+r_{ab|f0}-r_{d0ab0|da00c|0e000|00f00|0f000}-r_{gg000|0ab00|a00cb|e0000|0f000}.
\end{aligned}
$$

□

Remark 3.2. *It is hard to derive the maximal and minimal ranks of the general solution to (1.2) if we do not know the values of* $m_i, n_i, (i=1,2,3,4,6)$.

Similarly, we can get the corresponding results on X and Y. The proof is omitted.

Theorem 3.6. *Let* $A\in\mathbb{H}^{m\times n}, B\in\mathbb{H}^{m\times p_1}, C\in\mathbb{H}^{m\times p_2}, D\in\mathbb{H}^{m\times p_3}, E\in\mathbb{H}^{q_1\times n}, F\in\mathbb{H}^{q_2\times n}$ *and* $G\in\mathbb{H}^{q_3\times n}$ *be given. Assume that equation (1.2) is consistent. Then,*

$$
\begin{aligned}
\max_{BXE+CYF+DZG=A} r(X)=\min\big\{&p_1,\ q_1,\ p_1+q_1+r_{a|f|g}-r_b-r_{e|f|g},\ p_1+q_1+r_{acd}-r_e-r_{bcd},\\
&p_1+q_1+r_{b0ad0|ba00c|0g000|00f00}-r_{gg|e0|0f}-r_{bd}-r_{bc},\ p_1+q_1+r_{ac|g0}-r_{bc}-r_{f|g},\\
&p_1+q_1+r_{ad|f0}-r_{e|f}-r_{bd},\ p_1+q_1+r_{ee00|0ad0|a00c|g000|0f00}-r_{db0|d0c}-r_{e|g}-r_{e|f}\big\},
\end{aligned}
$$

$$
\begin{aligned}
&\min_{BXE+CYF+DZG=A} r\left(X\right)\\
=&r_{b0ad0|ba00c|0g000|00f00}+r_{ee00|0ad0|a00c|g000|0f00}+r_{acd}+r_{a|f|g}+r_{bd}+r_{e|g}+r_{cd}+r_{f|g}\\
&-r_{b0ad00|ba00cd|0g0000|00f000}-r_{ee00|0ad0|a00c|g000|0f00|0g00}\\
&+\max\left\{r_{ac|g0}-r_{b0ad0|ba00c|0g000|00f00|00g00}-r_{ee000|0ad0c|a00c0|g0000|0f000},\right.\\
&\qquad\left.r_{ad|f0}-r_{b0ad0|ba00c|0g000|00f00|0f000}-r_{ee000|0ad00|a00cd|g0000|0f000}\right\}.
\end{aligned}
$$

Theorem 3.7. *Let $A\in\mathbb{H}^{m\times n},B\in\mathbb{H}^{m\times p_1},C\in\mathbb{H}^{m\times p_2},D\in\mathbb{H}^{m\times p_3},E\in\mathbb{H}^{q_1\times n},F\in\mathbb{H}^{q_2\times n}$ and $G\in\mathbb{H}^{q_3\times n}$ be given. Assume that equation (1.2) is consistent. Then,*

$$
\begin{aligned}
\max_{BXE+CYF+DZG=A} r\left(Y\right)=\min\big\{&p_2,\ q_2,\ p_2+q_2+r_{a|e|g}-r_c-r_{e|f|g},\ p_2+q_2+r_{abd}-r_f-r_{bcd},\\
&p_2+q_2+r_{c0ab0|ca00d|0e000|00g00}-r_{gg|e0|0f}-r_{bc}-r_{cd},\ p_2+q_2+r_{ad|e0}-r_{cd}-r_{e|g},\\
&p_2+q_2+r_{ab|g0}-r_{f|g}-r_{bc},\ p_2+q_2+r_{ff00|0ab0|a00d|e000|0g00}-r_{db0|d0c}-r_{e|f}-r_{f|g}\big\},
\end{aligned}
$$

$$
\begin{aligned}
&\min_{BXE+CYF+DZG=A} r\left(Y\right)\\
=&r_{c0ab0|ca00d|0e000|00g00}+r_{ff00|0ab0|a00d|e000|0g00}+r_{abd}+r_{a|e|g}+r_{bc}+r_{e|f}+r_{bd}+r_{e|g}\\
&-r_{c0ab00|ca00db|0e0000|00g000}-r_{ff00|0ab0|a00d|e000|0g00|0e00}\\
&+\max\left\{r_{ad|e0}-r_{c0ab0|ca00d|0e000|00g00|00e00}-r_{ff000|0ab0d|a00d0|e0000|0g000},\right.\\
&\qquad\left.r_{ab|g0}-r_{c0ab0|ca00d|0e000|00g00|0g000}-r_{ff000|0ab00|a00db|e0000|0g000}\right\}.
\end{aligned}
$$

3.4. **Some solvability conditions and the general solution to (1.3).** In this section, the simultaneous decomposition of (1.1) will be used to solve the real quaternion matrix equation (1.3).

Theorem 3.8. *Let $A\in\mathbb{H}^{m\times n},B\in\mathbb{H}^{m\times p_1},C\in\mathbb{H}^{m\times p_2},D\in\mathbb{H}^{m\times p_3},E\in\mathbb{H}^{q_1\times n},F\in\mathbb{H}^{q_2\times n}$ and $G\in\mathbb{H}^{q_3\times n}$ be given. Then the equation (1.3) is consistent if and only if*

$$
r_{abcd|e000|f000|g000}=r_{bcd}+r_{e|f|g},\ A_{89}=0,\ A_{98}=0,\ A_{i,10}=0,\ A_{10,i}=0,\ (i=6,7,8,9). \tag{3.26}
$$

In this case, the general solution to (1.3) can be expressed as

$$
X=T_1^{-1}\widehat{X}Q,\quad W=P\widehat{W}V_1^{-1},\quad Y=T_2^{-1}\widehat{Y}V_2^{-1},\quad Z=T_3^{-1}\widehat{Z}V_3^{-1},
$$

where

$$\widehat{X} = \begin{array}{c} \\ m_1 \\ m_2 \\ m_3 \\ m_4 \\ m_5 \\ p_1 - r_b \end{array} \begin{array}{c} \begin{array}{ccccccccc} n_1 & n_2 & n_3 & n_4 & n_5 & n_4 & n_6 & n_7 & n_8 & n - r_{e|f|g} \end{array} \\ \begin{pmatrix} X_{11} & X_{12} & X_{13} & X_{14} & X_{15} & X_{16} & X_{17} & X_{18} & X_{19} & A_{1,10} \\ X_{21} & X_{22} & X_{23} & X_{24} & X_{25} & X_{26} & X_{27} & X_{28} & A_{29} & A_{2,10} \\ X_{31} & X_{32} & X_{33} & X_{34} & X_{35} & X_{36} & X_{37} & A_{38} & X_{39} & A_{3,10} \\ X_{41} & X_{42} & X_{43} & X_{44} & X_{45} & X_{46} & X_{47} & A_{48} & A_{49} - A_{69} & A_{4,10} \\ X_{51} & X_{52} & X_{53} & X_{54} & X_{55} & A_{56} & A_{57} & A_{58} & A_{59} & A_{5,10} \\ X_{61} & X_{62} & X_{63} & X_{64} & X_{65} & X_{66} & X_{67} & X_{68} & X_{69} & X_{6,10} \end{pmatrix} \end{array}, \tag{3.27}$$

$$\widehat{W} = \begin{array}{c} \\ m_1 \\ m_2 \\ m_3 \\ m_4 \\ m_5 \\ m_4 \\ m_6 \\ m_7 \\ m_8 \\ m - r_{bcd} \end{array} \begin{array}{c} \begin{array}{cccccc} n_1 & n_2 & n_3 & n_4 & n_5 & q_2 - r_e \end{array} \\ \begin{pmatrix} W_{11} & W_{12} & W_{13} & W_{14} & A_{15} - X_{15} & W_{16} \\ W_{21} & W_{22} & A_{23} - X_{23} & A_{24} - X_{24} & A_{25} - X_{25} & W_{26} \\ W_{31} & A_{32} - X_{32} & W_{33} & W_{34} & A_{35} - X_{35} & W_{36} \\ W_{41} & A_{42} - X_{42} & A_{43} - A_{63} + W_{63} - X_{43} & A_{44} - A_{46} + X_{46} - X_{44} & A_{45} - X_{45} & W_{46} \\ A_{51} - X_{51} & A_{52} - X_{52} & A_{53} - X_{53} & A_{54} - X_{54} & A_{55} - X_{55} & W_{56} \\ W_{61} & W_{62} & W_{63} & A_{64} - A_{46} + X_{46} & A_{65} & W_{66} \\ W_{71} & W_{72} & W_{73} & W_{74} & A_{75} & W_{76} \\ W_{81} & W_{82} & A_{83} & A_{84} & A_{85} & W_{86} \\ W_{91} & A_{92} & W_{93} & A_{94} - A_{96} & A_{95} & W_{96} \\ A_{10,1} & A_{10,2} & A_{10,3} & A_{10,4} & A_{10,5} & W_{10,6} \end{pmatrix} \end{array}, \tag{3.28}$$

$$\widehat{Y} = \begin{array}{c} \\ m_4 \\ m_6 \\ m_7 \\ m_1 \\ m_2 \\ p_2 - r_c \end{array} \begin{array}{c} \begin{array}{cccccc} n_4 & n_6 & n_7 & n_1 & n_2 & q_2 - r_f \end{array} \\ \begin{pmatrix} A_{66} - A_{64} + X_{46} & A_{67} - A_{47} + X_{47} & A_{68} & A_{61} - A_{41} + X_{41} + W_{41} - W_{61} & A_{62} - W_{62} & Y_{16} \\ A_{76} - A_{74} + W_{74} & Y_{22} & A_{78} & Y_{24} & A_{72} - W_{72} & Y_{26} \\ A_{86} & A_{87} & A_{88} & A_{81} - W_{81} & A_{82} - W_{82} & Y_{36} \\ A_{16} - A_{14} + X_{14} - X_{16} - W_{14} & Y_{42} & A_{18} - X_{18} & Y_{44} & A_{12} - (X_{12} + W_{12}) & Y_{46} \\ A_{26} - X_{26} & A_{27} - X_{27} & A_{28} - X_{28} & A_{21} - (X_{21} + W_{21}) & A_{22} - (X_{22} + W_{22}) & Y_{56} \\ Y_{61} & Y_{62} & Y_{63} & Y_{64} & Y_{65} & Y_{66} \end{pmatrix} \end{array}, \tag{3.29}$$

$\widehat{Z} =$

$$\begin{array}{c} \\ m_8 \\ m_4 \\ m_6 \\ m_3 \\ m_1 \\ p_3-r_d \end{array} \begin{array}{c} \begin{array}{cccccc} n_8 & n_4 & n_6 & n_3 & n_1 & q_3-r_g \end{array} \\ \begin{pmatrix} A_{99} & A_{96} & A_{97} & A_{93}-W_{93} & A_{91}-W_{91} & Z_{16} \\ A_{69} & A_{46}-X_{46} & A_{47}-X_{47} & A_{63}-W_{63} & A_{41}-(X_{41}+W_{41}) & Z_{26} \\ A_{79} & A_{74}-W_{74} & A_{77}-Y_{22} & A_{73}-W_{73} & A_{71}-(Y_{24}+W_{71}) & Z_{36} \\ A_{39}-X_{39} & A_{34}-(X_{34}+W_{34}) & A_{37}-X_{37} & A_{33}-(X_{33}+W_{33}) & A_{31}-(X_{31}+W_{31}) & Z_{46} \\ A_{19}-X_{19} & A_{14}-(X_{14}+W_{14}) & A_{17}-(X_{17}+Y_{42}) & A_{13}-(X_{13}+W_{13}) & Z_{55} & Z_{56} \\ Z_{61} & Z_{62} & Z_{63} & Z_{64} & Z_{65} & Z_{66} \end{pmatrix} \end{array}, \tag{3.30}$$

P, Q, A_{ij}, T_i, V_i are defined in Theorem 2.2, the remaining X_{ij}, Y_{ij}, Z_{ij} in (3.27)-(3.30) are arbitrary matrices over $\mathbb{H}$ with appropriate sizes.

Proof. From Theorem 2.2, we know that rewrite matrix equation (1.3) is consistent is equivalent to that the following matrix equation

$$P(S_B T_1 X Q^{-1} + P^{-1} W V_1 S_E + S_C T_2 Y V_2 S_F + S_D T_3 Z V_3 S_G)Q = P S_A Q.$$

Because P, Q are nonsingular, the matrix equation (1.3) can be written as

$$S_B T_1 X Q^{-1} + P^{-1} W V_1 S_E + S_C T_2 Y V_2 S_F + S_D T_3 Z V_3 S_G = S_A. \tag{3.31}$$

Let the matrices

$$\widehat{X} = T_1 X Q^{-1} = \begin{pmatrix} X_{11} & \cdots & X_{1,11} \\ \vdots & \ddots & \vdots \\ X_{61} & \cdots & X_{6,11} \end{pmatrix}, \ \widehat{W} = P^{-1} W V_1 = \begin{pmatrix} W_{11} & \cdots & W_{1,6} \\ \vdots & \ddots & \vdots \\ W_{11,1} & \cdots & W_{11,6} \end{pmatrix}, \tag{3.32}$$

$$\widehat{Y} = T_2 Y V_2 = \begin{pmatrix} Y_{11} & \cdots & Y_{16} \\ \vdots & \ddots & \vdots \\ Y_{61} & \cdots & Y_{66} \end{pmatrix}, \ \widehat{Z} = T_3 Z V_3 = \begin{pmatrix} Z_{11} & \cdots & Z_{16} \\ \vdots & \ddots & \vdots \\ Z_{61} & \cdots & Z_{66} \end{pmatrix}, \tag{3.33}$$

be partitioned in accordance with (3.31). Substituting (3.32) and (3.33) into (3.31) yields

$$S_A - S_B \widehat{X} - \widehat{W} S_E - S_C \widehat{Y} S_F - S_D \widehat{Z} S_G \triangleq \begin{pmatrix} \Omega_{11} & \Omega_{12} \\ \Omega_{21} & \Omega_{22} \end{pmatrix} = 0, \tag{3.34}$$

where

$\Omega_{11} =$

$$\begin{pmatrix} A_{11}-(X_{11}+W_{11}+Y_{44}+Z_{55}) & A_{12}-(X_{12}+W_{12}+Y_{45}) & A_{13}-(X_{13}+W_{13}+Z_{54}) & A_{14}-(X_{14}+W_{14}+Z_{52}) & A_{15}-(X_{15}+W_{15}) \\ A_{21}-(X_{21}+W_{21}+Y_{54}) & A_{22}-(X_{22}+W_{22}+Y_{55}) & A_{23}-(X_{23}+W_{23}) & A_{24}-(X_{24}+W_{24}) & A_{25}-(X_{25}+W_{25}) \\ A_{31}-(X_{31}+W_{31}+Z_{45}) & A_{32}-(X_{32}+W_{32}) & A_{33}-(X_{33}+W_{33}+Z_{44}) & A_{34}-(X_{34}+W_{34}+Z_{42}) & A_{35}-(X_{35}+W_{35}) \\ A_{41}-(X_{41}+W_{41}+Z_{25}) & A_{42}-(X_{42}+W_{42}) & A_{43}-(X_{43}+W_{43}+Z_{24}) & A_{44}-(X_{44}+W_{44}+Z_{22}) & A_{45}-(X_{45}+W_{45}) \\ A_{51}-(X_{51}+W_{51}) & A_{52}-(X_{52}+W_{52}) & A_{53}-(X_{53}+W_{53}) & A_{54}-(X_{54}+W_{54}) & A_{55}-(X_{55}+W_{55}) \end{pmatrix}, \tag{3.35}$$

$$\Omega_{12}=\begin{pmatrix} A_{16}-(Y_{41}+Z_{52}+X_{16}), & A_{17}-(Y_{42}+Z_{53}+X_{17}), & A_{18}-(Y_{43}+X_{18}), & A_{19}-(Z_{51}+X_{19}), & A_{1,10}-X_{1,10}, & -X_{1,11} \\ A_{26}-(Y_{51}+X_{26}), & A_{27}-(Y_{52}+X_{27}), & A_{28}-(Y_{53}+X_{28}), & A_{29}-X_{29}, & A_{2,10}-X_{2,10}, & -X_{2,11} \\ A_{36}-(Z_{42}+X_{36}), & A_{37}-(Z_{43}+X_{37}), & A_{38}-X_{38}, & A_{39}-(Z_{41}+X_{39}), & A_{3,10}-X_{3,10}, & -X_{3,11} \\ A_{46}-(Z_{22}+X_{46}), & A_{47}-(Z_{23}+X_{47}), & A_{48}-X_{48}, & A_{49}-(Z_{21}+X_{49}), & A_{4,10}-X_{4,10}, & -X_{4,11} \\ A_{56}-X_{56}, & A_{57}-X_{57} & A_{58}-X_{58}, & A_{59}-X_{59}, & A_{5,10}-X_{5,10}, & -X_{5,11} \end{pmatrix}, \tag{3.36}$$

$$\Omega_{21}=\begin{pmatrix} A_{61}-(Y_{14}+W_{61}+Z_{25}) & A_{62}-(Y_{15}+W_{62}) & A_{63}-(Z_{24}+W_{63}) & A_{64}-(Z_{22}+W_{64}) & A_{65}-W_{65} \\ A_{71}-(Y_{24}+W_{71}+Z_{35}) & A_{72}-(Y_{25}+W_{72}) & A_{73}-(Z_{34}+W_{73}) & A_{74}-(Z_{32}+W_{74}) & A_{75}-W_{75} \\ A_{81}-(Y_{34}+W_{81}) & A_{82}-(Y_{35}+W_{82}) & A_{83}-W_{83} & A_{84}-W_{84} & A_{85}-W_{85} \\ A_{91}-(Z_{15}+W_{91}) & A_{92}-W_{92} & A_{93}-(Z_{14}+W_{93}) & A_{94}-(Z_{12}+W_{94}) & A_{95}-W_{95} \\ A_{10,1}-W_{10,1} & A_{10,2}-W_{10,2} & A_{10,3}-W_{10,3} & A_{10,4}-W_{10,4} & A_{10,5}-W_{10,5} \\ -W_{11,1} & -W_{11,2} & -W_{11,3} & -W_{11,4} & -W_{11,5} \end{pmatrix}, \tag{3.37}$$

$$\Omega_{22}=\begin{pmatrix} A_{66}-(Y_{11}+Z_{22}) & A_{67}-(Y_{12}+Z_{23}) & A_{68}-Y_{13} & A_{69}-Z_{21} & A_{6,10} & 0 \\ A_{76}-(Y_{21}+Z_{32}) & A_{77}-(Y_{22}+Z_{33}) & A_{78}-Y_{23} & A_{79}-Z_{31} & A_{7,10} & 0 \\ A_{86}-Y_{31} & A_{87}-Y_{32} & A_{88}-Y_{33} & A_{89} & A_{8,10} & 0 \\ A_{96}-Z_{12} & A_{97}-Z_{13} & A_{98} & A_{99}-Z_{11} & A_{9,10} & 0 \\ A_{10,6} & A_{10,7} & A_{10,8} & A_{10,9} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & I_t \end{pmatrix}. \tag{3.38}$$

If the equation (1.3) has a solution (X,W,Y,Z), then by (3.34), we have that the equalities in (3.26) hold, and

$$\Omega_{11}=0,\Omega_{12}=0,\Omega_{21}=0,\Omega_{22}=0.$$

Conversely, assume that the equalities in (3.26) hold, then by (2.3)-(2.5) and (3.35)-(3.38), it can be verified that the matrices have the forms of (3.27)-(3.30) is a solution of (3.31), i.e., (1.2).

□

Remark 3.3. *The presented expressions of X and W are more useful than the expressions found by Wang and He* [21]*, since the latter can not be used to discuss the range of ranks of X and W to (1.3).*

3.5. **The range of ranks of the general solution to (1.3).** We in this section consider the range of ranks of the general solution to (1.3).

Theorem 3.9. *Let $A\in\mathbb{H}^{m\times n},B\in\mathbb{H}^{m\times p_1},C\in\mathbb{H}^{m\times p_2},D\in\mathbb{H}^{m\times p_3},E\in\mathbb{H}^{q_1\times n},F\in\mathbb{H}^{q_2\times n}$ and $G\in\mathbb{H}^{q_3\times n}$ be given. Assume that equation (1.3) is consistent. Then,*

$$\begin{aligned}\max_{BX+WE+CYF+DZG=A} r(X)=\min\big\{&p_1,\ n,\ p_1+r_{acd|e00}-r_{bcd},\ p_1+r_{a|e|f|g}-r_b,\\ &p_1+r_{b0ad0|ba00c|0g000|00f00|0e000|00e00}-r_{e|f|g}-r_{bc}-r_{bd},\\ &p_1+r_{ac|e0|g0}-r_{bc},\ p_1+r_{ad|e0|f0}-r_{bd}\big\},\end{aligned}$$

$$
\begin{aligned}
&\min_{BX+WE+CYF+DZG=A} r\left(X\right)\\
=&r_{a|e|f|g}+r_{acd|e00}-r_e-r_{b0ad00|ba00cd|0g0000|00f000|0e0000|00e000}\\
&+r_{bd}+r_{cd}+r_{b0ad0|ba00c|0g000|00f00|0e000|00e00}+r_{e|f|g}\\
&+\min\big\{r_{ac|e0|g0}-r_{b0ad0|ba00c|0g000|00f00|00g00|0e000|00e00}-r_{ad0c|a0c0|e000|f000|g000},\\
&\qquad r_{ad|e0|f0}-r_{b0ad0|ba00c|0g000|00f00|0e000|00e00|0f000}-r_{ad00|a0cd|e000|f000|g000}\big\}.
\end{aligned}
$$

Proof. It follows from Theorem 3.8 that the expression of X in (1.3) can be expressed as $X = T_1^{-1}\widehat{X}Q$, where $\widehat{X}$ is given in (3.27). Clearly, $r(X) = r(T_1^{-1}\widehat{X}Q) = r(\widehat{X})$. Now we consider the maximal and minimal ranks of $\widehat{X}$. Applying Lemma **??** to the variable matrices $\begin{pmatrix} X_{11} & \cdots & X_{15} \\ \vdots & \ddots & \vdots \\ X_{51} & \cdots & X_{55} \end{pmatrix}$ and $(X_{66},\ X_{67},\ X_{68},\ X_{69},\ X_{6,10})$ of $\widehat{X}$, we obtain

$$
\begin{aligned}
&\max_{\begin{pmatrix} X_{11} & \cdots & X_{15} \\ \vdots & \ddots & \vdots \\ X_{51} & \cdots & X_{55} \end{pmatrix},X_{6i},(i=6,\ldots,10)} r(\widehat{X})\\
&=\min\left\{p_1,n,p_1-r_b+r_e+r(\Phi_1),r(X_{61},X_{62},X_{63},X_{64},X_{65})+r_b+n_4+n_6+n_7+n_8+n-r_{e|f|g}\right\},
\end{aligned}
$$

$$
\min_{\begin{pmatrix} X_{11} & \cdots & X_{15} \\ \vdots & \ddots & \vdots \\ X_{51} & \cdots & X_{55} \end{pmatrix},X_{6i},(i=6,\ldots,10)} r(\widehat{X})=\max\left\{r(\Phi_1),\ r(X_{61},X_{62},X_{63},X_{64},X_{65})\right\},
$$

where

$$
\Phi_1=\begin{array}{c} \\ m_1 \\ m_2 \\ m_3 \\ m_4 \\ m_5 \end{array}\begin{array}{c} \begin{array}{ccccc} n_4 & n_6 & n_7 & n_8 & n-r_{e|f|g} \end{array} \\ \begin{pmatrix} X_{16} & X_{17} & X_{18} & X_{19} & A_{1,10} \\ X_{26} & X_{27} & X_{28} & A_{29} & A_{2,10} \\ X_{36} & X_{37} & A_{38} & X_{39} & A_{3,10} \\ X_{46} & X_{47} & A_{48} & A_{49}-A_{69} & A_{4,10} \\ A_{56} & A_{57} & A_{58} & A_{59} & A_{5,10} \end{pmatrix} \end{array}.
$$

Note that

$$
\max\left\{r(X_{61},X_{62},X_{63},X_{64},X_{65})\right\}=\min\left\{p_1-r_b,\ r_b\right\},\ \min\left\{r(X_{61},X_{62},X_{63},X_{64},X_{65})\right\}=0.
$$

Hence, we obtain

$$
\max_{\begin{pmatrix} X_{11} & \cdots & X_{15} \\ \vdots & \ddots & \vdots \\ X_{51} & \cdots & X_{55} \end{pmatrix},X_{6,i},(i=1,\ldots,10)} r(\widehat{X})=\min\left\{p_1,n,p_1-r_b+r_e+r(\Phi_1)\right\},
$$

$$
\min_{\begin{pmatrix} X_{11} & \cdots & X_{15} \\ \vdots & \ddots & \vdots \\ X_{51} & \cdots & X_{55} \end{pmatrix},X_{6,i},(i=1,\ldots,10)} r(\widehat{X})=r(\Phi_1).
$$

Applying Lemma 3.3 to the variable matrices $\begin{pmatrix} X_{16} & X_{17} \\ X_{26} & X_{27} \\ X_{36} & X_{37} \\ X_{46} & X_{47} \end{pmatrix}$ of Φ_1, we obtain

$$\max_{\begin{pmatrix} X_{16} & X_{17} \\ X_{26} & X_{27} \\ X_{36} & X_{37} \\ X_{46} & X_{47} \end{pmatrix}} r\left(\Phi_1\right) =$$

$$\min\left\{m_1+m_2+m_3+m_4+r\left(A_{56},\quad A_{57},\quad A_{58},\quad A_{59},\quad A_{5,10}\right), n_4+n_6+r(\Phi_2)\right\},$$

$$\min_{\begin{pmatrix} X_{16} & X_{17} \\ X_{26} & X_{27} \\ X_{36} & X_{37} \\ X_{46} & X_{47} \end{pmatrix}} r\left(\Phi_1\right) = r(\Phi_2)+r\left(A_{56},\quad A_{57},\quad A_{58},\quad A_{59},\quad A_{5,10}\right)-r\left(A_{58},\quad A_{59},\quad A_{5,10}\right),$$

where

$$\Phi_2 = \begin{array}{c} \\ m_1 \\ m_2 \\ m_3 \\ m_4 \\ m_5 \end{array} \begin{array}{c} \begin{array}{ccc} n_7 & n_8 & n-r_{e|f|g} \end{array} \\ \begin{pmatrix} X_{18} & X_{19} & A_{1,10} \\ X_{28} & A_{29} & A_{2,10} \\ A_{38} & X_{39} & A_{3,10} \\ A_{48} & A_{49}-A_{69} & A_{4,10} \\ A_{58} & A_{59} & A_{5,10} \end{pmatrix} \end{array}.$$

Applying Lemma 3.3 to the variable matrices $(X_{18},\ X_{19})$ of Φ_2, we obtain

$$\max_{(X_{18},\ X_{19})} r\left(\Phi_2\right) = \min\left\{n_7+n_8+r\begin{pmatrix} A_{1,10} \\ A_{2,10} \\ A_{3,10} \\ A_{4,10} \\ A_{5,10} \end{pmatrix}, m_1+r(\Phi_3)\right\},$$

$$\min_{(X_{18},\ X_{19})} r\left(\Phi_2\right) = r(\Phi_3)+r\begin{pmatrix} A_{1,10} \\ A_{2,10} \\ A_{3,10} \\ A_{4,10} \\ A_{5,10} \end{pmatrix} - r\begin{pmatrix} A_{2,10} \\ A_{3,10} \\ A_{4,10} \\ A_{5,10} \end{pmatrix},$$

where

$$\Phi_3 = \begin{array}{c} \\ m_2 \\ m_3 \\ m_4 \\ m_5 \end{array} \begin{array}{c} \begin{array}{ccc} n_7 & n_8 & n-r_{e|f|g} \end{array} \\ \begin{pmatrix} X_{28} & A_{29} & A_{2,10} \\ A_{38} & X_{39} & A_{3,10} \\ A_{48} & A_{49}-A_{69} & A_{4,10} \\ A_{58} & A_{59} & A_{5,10} \end{pmatrix} \end{array}.$$

Applying Lemma 3.1 to the variable matrices X_{28} and X_{39} of Φ_3, we obtain

$$\begin{aligned}
&\max_{X_{28},X_{39}} r\left(\Phi_3\right)\\
=&\min\left\{m_2+m_3+r\begin{pmatrix}A_{48} & A_{49}-A_{69} & A_{4,10}\\ A_{58} & A_{59} & A_{5,10}\end{pmatrix}, m_2+n_8+r\begin{pmatrix}A_{38} & A_{3,10}\\ A_{48} & A_{4,10}\\ A_{58} & A_{5,10}\end{pmatrix},\right.\\
&\left. m_3+n_7+r\begin{pmatrix}A_{29} & A_{2,10}\\ A_{49}-A_{69} & A_{4,10}\\ A_{59} & A_{5,10}\end{pmatrix}, n_7+n_8+r\begin{pmatrix}A_{2,10}\\ A_{3,10}\\ A_{4,10}\\ A_{5,10}\end{pmatrix}\right\},
\end{aligned}$$

$$\begin{aligned}
\min_{X_{28},X_{39}} r\left(\Phi_3\right)=&r\begin{pmatrix}A_{48} & A_{49}-A_{69} & A_{4,10}\\ A_{58} & A_{59} & A_{5,10}\end{pmatrix}+r\begin{pmatrix}A_{2,10}\\ A_{3,10}\\ A_{4,10}\\ A_{5,10}\end{pmatrix}\\
&+\max\left\{r\begin{pmatrix}A_{38} & A_{3,10}\\ A_{48} & A_{4,10}\\ A_{58} & A_{5,10}\end{pmatrix}-r\begin{pmatrix}A_{48} & A_{4,10}\\ A_{58} & A_{5,10}\end{pmatrix}-r\begin{pmatrix}A_{3,10}\\ A_{4,10}\\ A_{5,10}\end{pmatrix},\right.\\
&\left. r\begin{pmatrix}A_{29} & A_{2,10}\\ A_{49}-A_{69} & A_{4,10}\\ A_{59} & A_{5,10}\end{pmatrix}-r\begin{pmatrix}A_{49}-A_{69} & A_{4,10}\\ A_{59} & A_{5,10}\end{pmatrix}-r\begin{pmatrix}A_{2,10}\\ A_{4,10}\\ A_{5,10}\end{pmatrix}\right\}.
\end{aligned}$$

Hence,

$$\max_{BX+WE+CYF+DZG=A} r\left(X\right)=\min\left\{p_1,n,t_1,t_2,t_3,t_4,t_5,t_6\right\},$$

$$\min_{BX+WE+CYF+DZG=A} r\left(X\right)=\max\left\{t_7,t_8\right\},$$

where

$$t_1=p_1-r_b+r_e+m_1+m_2+m_3+m_4+r\left(A_{56},\ \ A_{57},\ \ A_{58},\ \ A_{59},\ \ A_{5,10}\right),$$

$$t_2=p_1-r_b+r_e+n_4+n_6+n_7+n_8+r\begin{pmatrix}A_{1,10}\\ A_{2,10}\\ A_{3,10}\\ A_{4,10}\\ A_{5,10}\end{pmatrix},$$

$$t_3=p_1-r_b+r_e+n_4+n_6+m_1+m_2+m_3+r\begin{pmatrix}A_{48} & A_{49}-A_{69} & A_{4,10}\\ A_{58} & A_{59} & A_{5,10}\end{pmatrix},$$

$$t_4 = p_1 - r_b + r_e + n_4 + n_6 + m_1 + m_2 + n_8 + r\begin{pmatrix} A_{38} & A_{3,10} \\ A_{48} & A_{4,10} \\ A_{58} & A_{5,10} \end{pmatrix},$$

$$t_5 = p_1 - r_b + r_e + n_4 + n_6 + m_1 + m_3 + n_7 + r\begin{pmatrix} A_{29} & A_{2,10} \\ A_{49} - A_{69} & A_{4,10} \\ A_{59} & A_{5,10} \end{pmatrix},$$

$$t_6 = p_1 - r_b + r_e + n_4 + n_6 + m_1 + n_7 + n_8 + r\begin{pmatrix} A_{2,10} \\ A_{3,10} \\ A_{4,10} \\ A_{5,10} \end{pmatrix} \geq t_2,$$

$$\begin{aligned} t_7 =& r\begin{pmatrix} A_{1,10} \\ A_{2,10} \\ A_{3,10} \\ A_{4,10} \\ A_{5,10} \end{pmatrix} + r\begin{pmatrix} A_{56}, & A_{57}, & A_{58}, & A_{59}, & A_{5,10} \end{pmatrix} + r\begin{pmatrix} A_{48} & A_{49} - A_{69} & A_{4,10} \\ A_{58} & A_{59} & A_{5,10} \end{pmatrix} \\ & - r(A_{58},\ A_{59},\ A_{5,10}) + r\begin{pmatrix} A_{38} & A_{3,10} \\ A_{48} & A_{4,10} \\ A_{58} & A_{5,10} \end{pmatrix} - r\begin{pmatrix} A_{48} & A_{4,10} \\ A_{58} & A_{5,10} \end{pmatrix} - r\begin{pmatrix} A_{3,10} \\ A_{4,10} \\ A_{5,10} \end{pmatrix}, \end{aligned}$$

$$\begin{aligned} t_8 =& r\begin{pmatrix} A_{1,10} \\ A_{2,10} \\ A_{3,10} \\ A_{4,10} \\ A_{5,10} \end{pmatrix} + r\begin{pmatrix} A_{56}, & A_{57}, & A_{58}, & A_{59}, & A_{5,10} \end{pmatrix} + r\begin{pmatrix} A_{48} & A_{49} - A_{69} & A_{4,10} \\ A_{58} & A_{59} & A_{5,10} \end{pmatrix} \\ & - r(A_{58},\ A_{59},\ A_{5,10}) + r\begin{pmatrix} A_{29} & A_{2,10} \\ A_{49} - A_{69} & A_{4,10} \\ A_{59} & A_{5,10} \end{pmatrix} - r\begin{pmatrix} A_{49} - A_{69} & A_{4,10} \\ A_{59} & A_{5,10} \end{pmatrix} - r\begin{pmatrix} A_{2,10} \\ A_{4,10} \\ A_{5,10} \end{pmatrix}. \end{aligned}$$

Now we pay attention to the ranks of the block matrices in t_i. Upon construction and computation, we obtain

$$\begin{aligned} r\begin{pmatrix} A_{56}, & A_{57}, & A_{58}, & A_{59}, & A_{5,10} \end{pmatrix} =& r\begin{pmatrix} S_A & S_C & S_D \\ S_E & 0 & 0 \end{pmatrix} - r(S_C,\ S_D) - r(S_E) \\ =& r_{acd|e00} - r_{cd} - r_e, \end{aligned} \tag{3.39}$$

$$r\begin{pmatrix} A_{1,10} \\ A_{2,10} \\ A_{3,10} \\ A_{4,10} \\ A_{5,10} \end{pmatrix} = r\begin{pmatrix} S_A \\ S_E \\ S_F \\ S_G \end{pmatrix} - r\begin{pmatrix} S_E \\ S_F \\ S_G \end{pmatrix} = r_{a|e|f|g} - r_{e|f|g}, \tag{3.40}$$

$$
\begin{aligned}
r\begin{pmatrix} A_{48} & A_{49}-A_{69} & A_{4,10} \\ A_{58} & A_{59} & A_{5,10} \end{pmatrix} =& r\begin{pmatrix} S_B & 0 & S_A & S_D & 0 \\ S_B & S_A & 0 & 0 & S_C \\ 0 & S_G & 0 & 0 & 0 \\ 0 & 0 & S_F & 0 & 0 \\ 0 & S_E & 0 & 0 & 0 \\ 0 & 0 & S_E & 0 & 0 \end{pmatrix} - r(S_C) - r(S_D) \\
& - r\begin{pmatrix} S_E \\ S_F \end{pmatrix} - r\begin{pmatrix} S_E \\ S_G \end{pmatrix} - m_2 - m_3 - m_4 - m_5 \\
=& r_{b0ad0|ba00c|0g000|00f00|0e000|00e00} \\
& - r_c - r_d - r_{e|f} - r_{e|g} - m_2 - m_3 - m_4 - m_5, \qquad (3.41)
\end{aligned}
$$

$$
r\begin{pmatrix} A_{38} & A_{3,10} \\ A_{48} & A_{4,10} \\ A_{58} & A_{5,10} \end{pmatrix} = r\begin{pmatrix} S_A & S_C \\ S_E & 0 \\ S_G & 0 \end{pmatrix} - r\begin{pmatrix} S_E \\ S_G \end{pmatrix} - r(S_C) = r_{ac|e0|g0} - r_{e|g} - r_c, \qquad (3.42)
$$

$$
r\begin{pmatrix} A_{29} & A_{2,10} \\ A_{49}-A_{69} & A_{4,10} \\ A_{59} & A_{5,10} \end{pmatrix} = r\begin{pmatrix} S_A & S_D \\ S_E & 0 \\ S_F & 0 \end{pmatrix} - r(S_D) - r\begin{pmatrix} S_E \\ S_F \end{pmatrix} = r_{ad|e0|f0} - r_d - r_{e|f}, \qquad (3.43)
$$

$$
\begin{aligned}
r\begin{pmatrix} A_{3,10} \\ A_{4,10} \\ A_{5,10} \end{pmatrix} =& r\begin{pmatrix} S_A & S_D & 0 & S_C \\ S_A & 0 & S_C & 0 \\ S_E & 0 & 0 & 0 \\ S_F & 0 & 0 & 0 \\ S_G & 0 & 0 & 0 \end{pmatrix} - r(S_C,\ S_D) - r(S_C) - r\begin{pmatrix} S_E \\ S_F \\ S_G \end{pmatrix} \\
=& r_{ad0c|a0c0|e000|f000|g000} - r_{cd} - r_c - r_{e|f|g}, \qquad (3.44)
\end{aligned}
$$

$$
\begin{aligned}
r\begin{pmatrix} A_{2,10} \\ A_{4,10} \\ A_{5,10} \end{pmatrix} =& r\begin{pmatrix} S_A & S_D & 0 & 0 \\ S_A & 0 & S_C & S_D \\ S_E & 0 & 0 & 0 \\ S_F & 0 & 0 & 0 \\ S_G & 0 & 0 & 0 \end{pmatrix} - r(S_C,\ S_D) - r(S_D) - r\begin{pmatrix} S_E \\ S_F \\ S_G \end{pmatrix} \\
=& r_{ad00|a0cd|e000|f000|g000} - r_{cd} - r_d - r_{e|f|g}, \qquad (3.45)
\end{aligned}
$$

$$
\begin{aligned}
r(A_{58},\ A_{59},\ A_{5,10}) =& r\begin{pmatrix} S_B & 0 & S_A & S_D & 0 & 0 \\ S_B & S_A & 0 & 0 & S_C & S_D \\ 0 & S_G & 0 & 0 & 0 & 0 \\ 0 & 0 & S_F & 0 & 0 & 0 \\ 0 & S_E & 0 & 0 & 0 & 0 \\ 0 & 0 & S_E & 0 & 0 & 0 \end{pmatrix} \\
& - r(S_B,\ S_D) - r(S_C,\ S_D) - r\begin{pmatrix} S_E \\ S_F \end{pmatrix} - r\begin{pmatrix} S_E \\ S_G \end{pmatrix} \\
=& r_{b0ad00|ba00cd|0g0000|00f000|0e0000|00e000} \\
& - r_{bd} - r_{cd} - r_{e|f} - r_{e|g}, \qquad (3.46)
\end{aligned}
$$

$$
\begin{aligned}
r\begin{pmatrix} A_{49} - A_{69} & A_{4,10} \\ A_{59} & A_{5,10} \end{pmatrix} =& r\begin{pmatrix} S_B & 0 & S_A & S_D & 0 \\ S_B & S_A & 0 & 0 & S_C \\ 0 & S_G & 0 & 0 & 0 \\ 0 & 0 & S_F & 0 & 0 \\ 0 & S_E & 0 & 0 & 0 \\ 0 & 0 & S_E & 0 & 0 \\ 0 & S_F & 0 & 0 & 0 \end{pmatrix} - r(S_C) - r(S_D) - r\begin{pmatrix} S_E \\ S_F \\ S_G \end{pmatrix} - r\begin{pmatrix} S_E \\ S_F \end{pmatrix} \\
& - m_2 - m_3 - m_4 - m_5 \\
=& r_{b0ad0|ba00c|0g000|00f00|0e000|00e00|0f000} \\
& - r_c - r_d - r_{e|f|g} - r_{e|f} - m_2 - m_3 - m_4 - m_5, \qquad (3.47)
\end{aligned}
$$

$$
\begin{aligned}
r\begin{pmatrix} A_{48} & A_{4,10} \\ A_{58} & A_{5,10} \end{pmatrix} =& r\begin{pmatrix} S_B & 0 & S_A & S_D & 0 \\ S_B & S_A & 0 & 0 & S_C \\ 0 & S_G & 0 & 0 & 0 \\ 0 & 0 & S_F & 0 & 0 \\ 0 & 0 & S_G & 0 & 0 \\ 0 & S_E & 0 & 0 & 0 \\ 0 & 0 & S_E & 0 & 0 \end{pmatrix} - r(S_C) - r(S_D) - r\begin{pmatrix} S_E \\ S_F \\ S_G \end{pmatrix} - r\begin{pmatrix} S_E \\ S_G \end{pmatrix} \\
& - m_3 - m_4 - m_5 \\
=& r_{b0ad0|ba00c|0g000|00f00|00g00|0e000|00e00} \\
& - r_c - r_d - r_{e|f|g} - r_{e|g} - m_2 - m_3 - m_4 - m_5. \qquad (3.48)
\end{aligned}
$$

Hence from (2.6)-(2.11) and (3.39)-(3.48), we deduce that

$$t_1 = p_1 - r_b + r_e + m_1 + m_2 + m_3 + m_4 + r_{acd|e00} - r_{cd} - r_e = p_1 + r_{acd|e00} - r_{bcd},$$

$$t_2 = p_1 - r_b + r_e + n_4 + n_6 + n_7 + n_8 + r_{a|e|f|g} - r_{e|f|g} = p_1 + r_{a|e|f|g} - r_b,$$

$$t_3 = p_1 + r_{b0ad0|ba00c|0g000|00f00|0e000|00e00} - r_{e|f|g} - r_{bc} - r_{bd},$$

$$t_4 = p_1 - r_b + r_e + n_4 + n_6 + m_1 + m_2 + n_8 + r_{ac|e0|g0} - r_{e|g} - r_c = p_1 + r_{ac|e0|g0} - r_{bc},$$

$$t_5 = p_1 - r_b + r_e + n_4 + n_6 + m_1 + m_3 + n_7 + r_{ad|e0|f0} - r_d - r_{e|f} = p_1 + r_{ad|e0|f0} - r_{bd},$$

$$\begin{aligned} t_7 =& r_{a|e|f|g} + r_{acd|e00} - r_e - r_{b0ad00|ba00cd|0g0000|00f000|0e0000|00e000} \\ &+ r_{bd} + r_{cd} + r_{b0ad0|ba00c|0g000|00f00|0e000|00e00} + r_{e|f|g} \\ &+ r_{ac|e0|g0} - r_{b0ad0|ba00c|0g000|00f00|00g00|0e000|00e00} - r_{ad0c|a0c0|e000|f000|g000}, \end{aligned}$$

$$\begin{aligned} t_8 =& r_{a|e|f|g} + r_{acd|e00} - r_e - r_{b0ad00|ba00cd|0g0000|00f000|0e0000|00e000} \\ &+ r_{bd} + r_{cd} + r_{b0ad0|ba00c|0g000|00f00|0e000|00e00} + r_{e|f|g} \\ &+ r_{ad|e0|f0} - r_{b0ad0|ba00c|0g000|00f00|0e000|00e00|0f000} - r_{ad00|a0cd|e000|f000|g000}. \end{aligned}$$

□

Similarly, we can get the corresponding results on W, Y, and Z.

Theorem 3.10. *Let $A \in \mathbb{H}^{m\times n}, B \in \mathbb{H}^{m\times p_1}, C \in \mathbb{H}^{m\times p_2}, D \in \mathbb{H}^{m\times p_3}, E \in \mathbb{H}^{q_1\times n}, F \in \mathbb{H}^{q_2\times n}$ and $G \in \mathbb{H}^{q_3\times n}$ be given. Assume that equation (1.3) is consistent. Then,*

$$\begin{aligned} \max_{BX+WE+CYF+DZG=A} r(W) =& \min\big\{q_1,\ m,\ q_1 + r_{ab|f0|g0} - r_{e|f|g},\ q_1 + r_{abcd} - r_e, \\ & q_1 + r_{ee0000|0ad0b0|a00c0b|g00000|0f0000} - r_{bcd} - r_{e|f} - r_{e|g}, \\ & q_1 + r_{abd|f00} - r_{e|f},\ q_1 + r_{abc|g00} - r_{e|g}\big\}, \end{aligned}$$

$$\begin{aligned} & \min_{BX+WE+CYF+DZG=A} r(W) \\ =& r_{abcd} + r_{ab|f0|g0} - r_b - r_{ee0000|0ad0b0|a00c0b|g00000|0f0000|0g0000} + r_{e|g} + r_{f|g} \\ &+ r_{ee0000|0ad0b0|a00c0b|g00000|0f0000} + r_{bcd} \\ &+ \min\big\{r_{abd|f00} - r_{ee00000|0ad00b0|a00cd0b|g000000|0f00000} - r_{aadcb|g0000|0f000|f0000}, \\ & \qquad r_{abc|g00} - r_{ee00000|0ad0b0c|a00c0b0|g000000|0f00000} - r_{aabcd|g0000|0f000|0g000}\big\}. \end{aligned}$$

Theorem 3.11. *Let $A \in \mathbb{H}^{m\times n}, B \in \mathbb{H}^{m\times p_1}, C \in \mathbb{H}^{m\times p_2}, D \in \mathbb{H}^{m\times p_3}, E \in \mathbb{H}^{q_1\times n}, F \in \mathbb{H}^{q_2\times n}$ and $G \in \mathbb{H}^{q_3\times n}$ be given. Assume that equation (1.3) is consistent. Then,*

$$\begin{aligned} & \max_{BX+WE+CYF+DZG=A} r(Y) = \\ & \min\big\{p_2,\ q_2,\ p_2 + q_2 + r_{ab|e0|g0} - r_{e|f|g} - r_{bc},\ p_2 + q_2 + r_{abd|e00} - r_{e|f} - r_{bcd}\big\}, \\ & \min_{BX+WE+CYF+DZG=A} r(Y) = r_{ab|e0|g0} + r_{abd|e00} - r_{abd|e00|g00} - r_b - r_e, \\ & \max_{BX+WE+CYF+DZG=A} r(Z) = \\ & \min\big\{p_3,\ q_3,\ p_3 + q_3 + r_{ab|e0|f0} - r_{e|f|g} - r_{bd},\ p_3 + q_3 + r_{abc|e00} - r_{e|g} - r_{bcd}\big\}, \\ & \min_{BX+WE+CYF+DZG=A} r(Z) = r_{ab|e0|f0} + r_{abc|e00} - r_{abc|e00|f00} - r_b - r_e. \end{aligned}$$

Remark 3.4. *All the results are true over octonion algebra.*

4. **Conclusion**

We have established the simultaneous decomposition of the general real quaternion matrix array (1.1). We have derived all the dimensions of identity matrices in the equivalence canonical form of the matrix array (1.1). Using the simultaneous decomposition of the general matrix array (1.1), we have presented necessary and sufficient conditions for the existence and the general solutions to the real matrix equations (1.2) and (1.3), respectively. Moreover, we have given the range of ranks of the general solutions to (1.2) and (1.3), respectively.

As a special case of the matrix array (1.1), we have derived all the dimensions of identity matrices in the equivalence canonical form of triple matrices with the same row or column numbers, which perfect the results in [19]. The presented expression of the general solution is more useful than the expression found in [19], since the latter can not be used to consider the maximal and minimal ranks of the general solution to (1.2). On the other hand, Wang and He [21] gave the range of ranks of Y and Z to (1.3), but the two present authors did not derive the range of ranks of X and W. We in this paper have solved this problem.

References

[1] N.L. Bihan, J. Mars, Singular value decomposition of quaternion matrices: A new tool for vector-sensor signal processing, *Signal Processing.* 84 (7) (2004) 1177–1199.

[2] J.C.K. Chou, Quaternion kinematic and dynamic differential equations, *IEEE Trans. Robot. Automat.* 8 (1992) 53–64.

[3] B. De Moor, H.Y. Zha, A tree of generalization of the ordinary singular value decomposition, *Linear Algebra Appl.* 147 (1991) 469–500.

[4] B. De Moor, G.H. Golub, The restricted singular value decomposition: properties and applications, *SIAM J. Matrix Anal. Appl.* 12 (3) (1991) 401–425.

[5] B. De Moor, Paul Van Dooren, Generalizations of the singular value and QR decompositions, *SIAM J. Matrix Anal. Appl.* 13 (4) (1992) 993–1014.

[6] D.L Chu, L. De Lathauwer, B. De Moor, On the computation of restricted singular value decomposition via cosine-sine decomposition, *SIAM J. Matrix Anal. Appl.* 22 (2) (2000) 550–601.

[7] D.L. Chu, Y.S. Hung, H.J. Woerdeman, Inertia and rank characterizations of some matrix expressions, *SIAM J. Matrix Anal. Appl.* 31 (2009) 1187–1226.

[8] N. Cohen, C.R. Johnson, L. Rodman, H.J. Woerdeman, Ranks of completions of partial matrices, *Oper. Theory: Adv. Appl.* 40 (1989) 165–185.

[9] Z.H. He, Q.W. Wang, A simultaneous decomposition of five real quaternion matrices with applications, http://arxiv.org/abs/1306.5074v1.

[10] T.W. Hungerford, Algebra, Springer-Verlag, New York, 1980.

[11] S. De Leo, G. Scolarici, Right eigenvalue equation in quaternionic quantum mechanics, *J. Phys. A.* 33 (2000) 2971–2995.

[12] C.C. Paige, M.A. Saunders, Towards a generalized singular value decomposition, *SIAM J. Number. Anal.* 18 (1981) 398–405.

[13] C.C. Paige, M.S. Wei, History and generality of the CS decomposition, *Linear Algebra Appl.* 208-209 (1994) 303–326.

[14] B. De Schutter, B. De Moor, The QR decomposition and the singular value decomposition in the symmetrized max-plus algebra revisited, *SIAM review.* 44(3) 2002 417-454.

[15] C.C. Took, D.P. Mandic, Augmented second-order statistics of quaternion random signals, *Signal Processing.* 91 (2011) 214–224.

[16] C.C. Took, D.P. Mandic, The quaternion LMS algorithm for adaptive filtering of hypercomplex real world processes, *IEEE Trans. Signal Process.* 57 (2009) 1316–1327.

[17] C.C. Took, D.P. Mandic, A quaternion widely linear adaptive filter, *IEEE Trans. Signal Process.* 58 (2010) 4427–4431.

[18] C.C. Took, D.P. Mandic, F. Zhang, On the unitary diagonalisation of a special class of quaternion matrices, *Appl. Math. Lett.* 24 (2011) 1806–1809.

[19] Q.W. Wang, J.W. van der Woude, S.W. Yu, An equivalence canonical form of a matrix triplet over an arbitrary division ring with applications, *Sci. China Math.* 54 (5)(2011) 907–924.

[20] Q.W. Wang, X. Zhang, J.W. van der Woude, A new simultaneous decomposition of a matrix quaternity over an arbitrary division ring with applications, *Comm. Algebra.* 40 (2012) 2309-2342.

[21] Q.W. Wang, Z.H. He, Some matrix equations with applications, *Linear Multilinear Algebra.* 60 (2012) 1327–1353.

[22] H.J. Woerdeman, The lower order of triangular operators and minimal rank extensions, *Integral Equations operator Theory.* 10 (1987) 859–879.

[23] H.J. Woerdeman, Minimal rank completions for block matrices, *Linear Algebra Appl.* 121 (1989) 105–122.

[24] H.J. Woerdeman, Minimal rank completions of partial banded matrices, *Linear and Multilinear Algebra.* 36 (1993) 59–69.

[25] H.Y. Zha, The restricted singular value decomposition of matrix triplets, *SIAM J. Matrix Anal. Appl.* 12 (1991) 172–194.

[26] F.Z. Zhang, Quaternions and matrices of quaternions, *Linear Algebra Appl.* 251 (1997) 21–57.